

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
7 October 2004 (07.10.2004)

PCT

(10) International Publication Number  
WO 2004/085385 A2

(51) International Patent Classification<sup>7</sup>: C07C 317/00

PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(21) International Application Number:  
PCT/US2004/008333

(22) International Filing Date: 18 March 2004 (18.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/456,268 20 March 2003 (20.03.2003) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

#### Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (for all designated States except US): SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): SHANKAR, Bandarpalle, B. [US/US]; 1124 Van Arsdale Drive, Branchburg, NJ 08853 (US). RIZVI, Razia, K. [US/US]; 22 Tomar Court, Bloomfield, NJ 07003 (US). KOZLOWSKI, Joseph, A. [US/US]; 1066 Stuart Road, Princeton, NJ 08540 (US). SHIH, Neng-Yang [US/US]; 4 Overlook Drive, Warren, NJ 07059 (US).

(74) Agent: BLINKA, Thomas, A.; Schering-Plough Corporation, Patent Department, K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,

(54) Title: CANNABINOID RECEPTOR LIGANDS



(57) Abstract: There are disclosed compounds of the formula (I) a pharmaceutically acceptable salt or solvate of the compound, which exhibit anti-inflammatory and immunomodulatory activity. Also disclosed are pharmaceutical compositions containing said compounds.

## CANNABINOID RECEPTOR LIGANDS

### CROSS-REFERENCE TO RELATED APPLICATION

This application claims the benefit of priority from U.S. Provisional Patent Application Serial No. 60/456,268, filed March 20, 2003.

### BACKGROUND OF THE INVENTION

#### Field of the Invention

The present invention relates to cannabinoid receptor ligands and, more particularly, to compounds that bind to cannabinoid (CB<sub>2</sub>) receptors. Compounds according to the present invention generally exhibit anti-inflammatory and immunomodulatory activity and can be useful in treating conditions characterized by inflammation and immunomodulatory irregularities. Examples of conditions which may be treated include, but are not limited to, rheumatoid arthritis, asthma, allergy, psoriasis, Crohn's disease, systemic lupus erythematosus, multiple sclerosis, diabetes, cancer, glaucoma, osteoporosis, renal ischemia, cerebral stroke, cerebral ischemia, and nephritis. The invention also relates to pharmaceutical compositions containing said compounds.


Cannabinoid receptors belong to the superfamily of G-protein coupled receptors. They are classified into the predominantly neuronal CB<sub>1</sub> receptors and the predominantly peripheral CB<sub>2</sub> receptors. While the effects of CB<sub>1</sub> receptors are principally associated with the central nervous system, CB<sub>2</sub> receptors are believed to have peripheral effects related to bronchial constriction, immunomodulation and inflammation. As such, a selective CB<sub>2</sub> receptor binding agent is expected to have therapeutic utility in the control of diseases associated with inflammation, immunomodulation and bronchial constriction such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, diabetes, osteoporosis, renal ischemia, cerebral stroke, cerebral ischemia, nephritis, inflammatory disorders of the lungs and gastrointestinal tract, and respiratory tract disorders such as reversible airway obstruction, chronic asthma and bronchitis (see, e.g., R.G. Pertwee, Curr. Med. Chem. 6(8), (1999), 635).

Various compounds have reportedly been developed which interact with CB<sub>2</sub> receptors and/or which have, *inter alia*, anti-inflammatory activity associated with

cannabinoid receptors. See, e.g., U.S. Pat. Nos. 5,338,753, 5,462,960, 5,532,237, 5,925,768, 5,948,777, 5,990,170, 6,013,648 and 6,017,919.

### SUMMARY OF THE INVENTION

5 In its many embodiments, the present invention provides compounds of formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein

n is 0 to 4;

10 p is 0 to 4;

q is 0 to 5;

X is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, haloalkoxy, haloalkyl, halogen, heteroalkyl, heteroaryl, -CF<sub>3</sub>, -CN, -C(O)N(R<sup>2</sup>)<sub>2</sub>, -C(O)OR<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub>, -NHC(O)R<sup>2</sup>,  
 15 -NR<sup>2</sup>C(O)OR<sup>2</sup>, -NR<sup>2</sup>C(O)N(R<sup>2</sup>)<sub>2</sub>, -NO<sub>2</sub>, -NC(=N-CN)NHR<sup>2</sup>, -OCF<sub>2</sub>H, -OCF<sub>3</sub>, -OH and -S(O<sub>2</sub>)N(R<sup>2</sup>)<sub>2</sub>, with the proviso that when n is 2, 3 or 4, the X moieties can be the same or different and are independently selected from the group listed above;

Y is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, halogen, heteroalkyl, heteroaryl,  
 20 -O-cycloalkyl, -CF<sub>3</sub>, -CN, -C(O)OR<sup>2</sup>, -C(O)R<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -OCF<sub>2</sub>H and -OH, with the proviso that:

when p is 2, 3 or 4, the Y moieties can be the same or different and are independently selected from the group listed above; or

when p is 2, the Y moieties can form a cyclic ring of 3 to 7 ring atoms of which  
 25 1-2 may be a heteroatom;

Z is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl, cycloalkenyl, halogen, heteroalkyl, heteroaryl, heterocyclyl, heterocyclenyl,

-O-cycloalkyl, -CF<sub>3</sub>, -CN, -C(O)OR<sup>2</sup>, -C(O)R<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -OCF<sub>2</sub>H, -OH and -S(O<sub>2</sub>)R<sup>2</sup>, with the proviso that when q is 2, 3, 4 or 5, the Z moieties can be the same or different and are independently selected from the group listed above;

R<sup>1</sup> is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl, heterocyclenyl, and -N(R<sup>2</sup>)<sub>2</sub>;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl, heterocyclenyl, and hydroxyalkyl;

L<sup>1</sup> is selected from the group consisting of a covalent bond, -C(F<sub>2</sub>)-, -CH(OR<sup>2</sup>)-, -C(O)-, -C(O)N(H)-, -C(=N-OR<sup>2</sup>)-, -C(=NR<sup>2</sup>)-, -C(=N-CN)-, -C(R<sup>2</sup>)<sub>2</sub>-, -N(R<sup>2</sup>)-, -N(H)C(O)-, -N(R<sup>2</sup>)S(O<sub>2</sub>)-, -O-, -OC(O)-, -C(O)O-, -S-, -S(O<sub>2</sub>)-, -S(O)- and -S(O<sub>2</sub>)N(R<sup>2</sup>)-, with the proviso that when L<sup>1</sup> is a covalent bond, M<sup>1</sup> is directly attached to the phenyl carbon marked A;

L<sup>2</sup> is selected from the group consisting of a covalent bond, -C(R<sup>2</sup>)<sub>2</sub>-, -C(=N-OR<sup>2</sup>)-, -C(O)-, -C(O)N(H)-, -C(O)O-, -OC(O)-, -N(H)C(O)-, -NHS(O<sub>2</sub>)-, -N(R<sup>2</sup>)-, -O-, -S-, -S(O)-, -S(O<sub>2</sub>)- and -S(O<sub>2</sub>)N(R<sup>2</sup>)-, with the proviso that when L<sup>2</sup> is a covalent bond, M<sup>1</sup> is directly attached to M<sup>2</sup>;

M<sup>1</sup> is aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl or heterocyclenyl; and

M<sup>2</sup> is alkyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl, heterocyclenyl, -C(O)R<sup>2</sup>, -C(O)OR<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub> or -S(O<sub>2</sub>)R<sup>2</sup>;

with the provisos that:

- i) when M<sup>2</sup> is -N(R<sup>2</sup>)<sub>2</sub>, L<sup>2</sup> is a covalent bond, -CH<sub>2</sub>-, -C(O)- or -S(O<sub>2</sub>)- and Z is absent;
- ii) when M<sup>2</sup> is -C(O)R<sup>2</sup> or -C(O)OR<sup>2</sup>, L<sup>2</sup> is a covalent bond, -CH<sub>2</sub>-, -NH- or -N(alkyl)- and Z is absent (i.e., q is 0);
- iii) when M<sup>2</sup> is -S(O<sub>2</sub>)R<sup>2</sup>, L<sup>2</sup> is a covalent bond, -CH<sub>2</sub>-, -NH- or -N(alkyl)- and Z is absent (i.e., q is 0); and
- iv) the two R<sup>2</sup> moieties of -N(R<sup>2</sup>)<sub>2</sub> and -C(R<sup>2</sup>)<sub>2</sub>- are the same or different and are independently selected, or the two R<sup>2</sup> of -N(R<sup>2</sup>)<sub>2</sub> are joined together and with the nitrogen to which they are attached to form a heterocyclic ring having 3 to 7 ring atoms optionally containing additionally one or more N or O atoms (such as, for example, morpholine, piperidine and piperazine) wherein said additional N can be optionally substituted with R<sup>2</sup>;

wherein each of said alkoxy, alkyl, aralkyl, aryl, cycloalkyl, cycloalkenyl, heteroaralkyl, heteroaryl, heterocyclyl, and heterocyclenyl in the definitions above can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkoxy,  $-\text{CF}_3$ ,  $-\text{CN}$ ,  $-\text{C}(\text{O})\text{N}(\text{R}^2)_2$ ,  $-\text{C}(\text{O})\text{OR}^2$ ,  $-\text{C}(\text{O})\text{R}^2$ ,  $-\text{NC}(\text{O})\text{R}^2$ ,  $-\text{NR}^2\text{C}(\text{O})\text{OR}^2$ ,  $-\text{NR}^2\text{C}(\text{O})\text{N}(\text{R}^2)_2$ ,  $-\text{NC}(=\text{N}-\text{CN})\text{NHR}^2$ ,  $-\text{NO}_2$ ,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_2\text{H}$ ,  $-\text{OCF}_3$ ,  $-\text{OH}$ ,  $-\text{S}(\text{O}_2)\text{R}^2$  and  $-\text{S}(\text{O}_2)\text{N}(\text{R}^2)_2$ .

The compounds of the present invention can be useful as cannabinoid receptor ligands. The compounds can have anti-inflammatory activity and/or immunomodulatory activity and can be useful in the treatment of various medical conditions including, e.g., cutaneous T-cell lymphoma, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, sepsis, shock, sarcoidosis, idiopathic pulmonary fibrosis, bronchopulmonary dysplasia, retinal disease, scleroderma, osteoporosis, renal ischemia, myocardial infarction, cerebral stroke, cerebral ischemia, nephritis, hepatitis, glomerulonephritis, cryptogenic fibrosing aveolitis, psoriasis, transplant rejection, atopic dermatitis, vasculitis, allergy, seasonal allergic rhinitis, Crohn's disease, inflammatory bowel disease, reversible airway obstruction, adult respiratory distress syndrome, asthma, chronic obstructive pulmonary disease (COPD) or bronchitis. It is contemplated that a compound of this invention can be useful in treating one or more of the diseases listed.

#### DETAILED DESCRIPTION

In one embodiment, the present invention discloses cannabinoid receptor ligands represented by structural formula I, or a pharmaceutically acceptable salt or solvate thereof, wherein the various moieties are described above.

In a preferred embodiment of compounds of formula I,

$\text{M}^1$  is aryl, heteroaryl or heterocyclyl;

and

$\text{M}^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl.

In another preferred embodiment, X is selected from the group consisting of hydrogen, alkoxy,  $-\text{CF}_3$ , haloalkoxy, halogen,  $-\text{OCF}_3$ ,  $-\text{OCF}_2\text{H}$  and  $-\text{OH}$ .

In another preferred embodiment, Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , cycloalkyl, halogen,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$ ,

-O-cycloalkyl and -OH.

In another preferred embodiment, Z is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , halogen, heterocyclyl,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$ , -O-cycloalkyl and -OH.

5 In another preferred embodiment,  $\text{R}^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-\text{N}(\text{R}^2)_2$ .

In another preferred embodiment,  $\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl and heterocyclyl.

10 In another preferred embodiment,  $\text{L}^1$  is selected from the group consisting of  $-\text{C}(\text{R}^2)_2-$ ,  $-\text{N}(\text{R}^2)\text{S}(\text{O}_2)-$ ,  $-\text{N}(\text{R}^2)-$ ,  $-\text{S}(\text{O}_2)-$  and  $-\text{S}(\text{O}_2)\text{N}(\text{R}^2)-$ .

In another preferred embodiment,  $\text{L}^2$  is selected from the group consisting of a covalent bond,  $-\text{C}(\text{R}^2)_2-$ ,  $-\text{N}(\text{R}^2)\text{S}(\text{O}_2)-$ ,  $-\text{N}(\text{R}^2)-$ ,  $-\text{S}(\text{O}_2)-$  and  $-\text{S}(\text{O}_2)\text{N}(\text{R}^2)-$ .

In another preferred embodiment,  $\text{M}^1$  is aryl, heteroaryl or heterocyclyl.

15 In another preferred embodiment,  $\text{M}^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl.

In an additional preferred embodiment, n is 1.

In an additional preferred embodiment, X is hydrogen and/or -OH.

20 In an additional preferred embodiment, Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , cycloalkyl, halogen,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$  and -OH.

In an additional preferred embodiment, Z is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , halogen, heterocyclyl,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$ , -O-cycloalkyl and -OH.

25 In an additional preferred embodiment,  $\text{R}^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-\text{N}(\text{R}^2)_2$ .

In an additional preferred embodiment,  $\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl and heterocyclyl.

In an additional preferred embodiment,  $\text{L}^1$  is selected from the group consisting of  $-\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^2)_2-$  and  $-\text{S}(\text{O}_2)-$ .

30 In an additional preferred embodiment,  $\text{L}^2$  is selected from the group consisting of a covalent bond,  $-\text{C}(\text{R}^2)_2-$ ,  $-\text{C}(\text{O})-$  and  $-\text{S}(\text{O}_2)-$ .

In an additional preferred embodiment,  $\text{M}^1$  is aryl, heteroaryl or heterocyclyl.

In an additional preferred embodiment,  $M^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl.

In a still additional preferred embodiment,  $n$  is 1.

In a still additional preferred embodiment,  $X$  is hydrogen and/or  $-OH$ .

5 In a still additional preferred embodiment,  $Y$  is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-C(O)OR^2$ , cycloalkyl,  $-CF_3$ , halogen,  $-OCF_3$  and  $-OH$ .

In a still additional preferred embodiment,  $Z$  is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ , halogen,  $-OCF_3$ ,  $-O$ -cycloalkyl and  $-OH$ .

10 In a still additional preferred embodiment,  $R^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-N(R^2)_2$ .

In a still additional preferred embodiment,  $R^2$  is selected from the group consisting of hydrogen, alkyl, aryl and cycloalkyl.

15 In a still additional preferred embodiment,  $L^1$  and  $L^2$  are the same or different and are independently  $-C(R^2)_2$ - or  $-S(O_2)$ -.

In a still additional preferred embodiment,  $M^1$  is aryl or heteroaryl.

In a still additional preferred embodiment,  $M^2$  is aryl, cycloalkyl, heteroaryl or heterocyclyl.

20 A preferred embodiment of compounds has the structure of formula I or a pharmaceutically acceptable salt or solvate thereof, wherein

$X$  is selected from the group consisting of hydrogen, alkoxy,  $-CF_3$ , haloalkoxy, halogen,  $-OCF_3$ ,  $-OCF_2H$  and  $-OH$ ;

$Y$  is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ ,  $-C(O)OR^2$ , cycloalkyl, halogen,  $-N(R^2)_2$ ,  $-OCF_3$ ,  $-O$ -cycloalkyl and  $-OH$ ;

25  $Z$  is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ ,  $-C(O)OR^2$ , halogen, heterocyclyl,  $-N(R^2)_2$ ,  $-OCF_3$ ,  $-O$ -cycloalkyl and  $-OH$ ;

$R^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-N(R^2)_2$ ;

30  $R^2$  is selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl and heterocyclyl;

$L^1$  is selected from the group consisting of  $-C(R^2)_2$ -,  $-N(R^2)$ -,  $-S(O_2)$ - and  $-S(O_2)N(R^2)$ -;

$L^2$  is selected from the group consisting of a covalent bond,  $-C(R^2)_2-$ ,  $-N(R^2)S(O_2)-$ ,  $-N(R^2)-$ ,  $-S(O_2)-$  and  $-S(O_2)N(R^2)-$ ;

$M^1$  is aryl, heteroaryl or heterocycloalkyl;

$M^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl;

5 with the proviso that:

the two  $R^2$  moieties of  $-N(R^2)_2$  and  $-C(R^2)_2-$  are the same or different and are independently selected, or wherein each  $R^2$  of  $-N(R^2)_2$  are joined together and with the nitrogen to which they are attached form a heterocyclic ring having 3 to 7 ring atoms;

wherein each of said alkoxy, alkyl, aralkyl, aryl, cycloalkyl, heteroaralkyl, heteroaryl and heterocyclyl in the definitions above can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen,  $-CF_3$ ,  $-CN$ ,  $-C(O)N(R^2)_2$ ,  $-C(O)OR^2$ ,  $-NR^2C(O)R^2$ ,  $-NR^2C(O)OR^2$ ,  $-NR^2C(O)N(R^2)_2$ ,  $-NC(=N-CN)NHR^2$ ,  $-NO_2$ ,  $-N(R^2)_2$ ,  $-OCF_2H$ ,  $-OCF_3$ ,  $-OH$ ,  $-S(O_2)R^2$  and  $-S(O_2)N(R^2)_2$ .

Another preferred embodiment of compounds has the structure of formula I or a pharmaceutically acceptable salt or solvate thereof, wherein

$n$  is 1;

$X$  is  $-OH$ ;

20  $Y$  is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ ,  $-C(O)OR^2$ , cycloalkyl, halogen,  $-N(R^2)_2$ ,  $-OCF_3$  and  $-OH$ ;

$Z$  is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ ,  $-C(O)OR^2$ , halogen, heterocyclyl,  $-N(R^2)_2$ ,  $-OCF_3$ ,  $-O$ -cycloalkyl and  $-OH$ ;

25  $R^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-N(R^2)_2$ ;

$R^2$  is selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl and heterocyclyl;

$L^1$  is selected from the group consisting of  $-C(O)-$ ,  $-C(R^2)_2-$  and  $-S(O_2)-$ ;

30  $L^2$  is selected from the group consisting of a covalent bond,  $-C(R^2)_2-$ ,  $-C(O)-$  and  $-S(O_2)-$ ;

$M^1$  is aryl, heteroaryl or heterocyclyl;

and

$M^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl.



Another preferred embodiment of compounds has the structure of formula I or a pharmaceutically acceptable salt or solvate thereof, wherein

n is 1;

X is -OH;

5 Y is selected from the group consisting of hydrogen, alkoxy, alkyl, -C(O)OR<sup>2</sup>, cycloalkyl, -CF<sub>3</sub>, halogen, -OCF<sub>3</sub> and -OH;

Z is selected from the group consisting of hydrogen, alkoxy, alkyl, -CF<sub>3</sub>, halogen, -OCF<sub>3</sub>, -O-cycloalkyl and -OH;

10 R<sup>1</sup> is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and -N(R<sup>2</sup>)<sub>2</sub>;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, aryl and cycloalkyl;


L<sup>1</sup> and L<sup>2</sup> are the same or different and are independently -C(R<sup>2</sup>)<sub>2</sub>- or -S(O<sub>2</sub>)-;

M<sup>1</sup> is aryl or heteroaryl;

and

15 M<sup>2</sup> is aryl, cycloalkyl, heteroaryl or heterocyclyl.

Another preferred embodiment of compounds has the structure of formula II:



20 or a pharmaceutically acceptable salt thereof, wherein

Y is selected from the group consisting of hydrogen, alkoxy, alkyl, -CF<sub>3</sub>, -CN, -C(O)OR<sup>2</sup>, cycloalkyl, halogen, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub> and -OH;

Z is selected from the group consisting of hydrogen, alkoxy, alkyl, -CF<sub>3</sub>, -C(O)OR<sup>2</sup>, halogen, heterocyclyl, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -O-cycloalkyl and -OH;

25 R<sup>1</sup> is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and -N(R<sup>2</sup>)<sub>2</sub>;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, aryl and cycloalkyl;

L<sup>1</sup> is -C(R<sup>2</sup>)<sub>2</sub>- or -S(O<sub>2</sub>)-;










$M^1$  is aryl, indolyl or furanyl;













and
















$M^2$  is aryl, cycloalkyl, heteroaryl or heterocyclyl.

- 5 Additionally preferred compounds of formula II above, or a pharmaceutically acceptable salt thereof, are shown below, wherein  $R^1$ ,  $L^1$ ,  $M^1$ ,  $M^2$ , Y and Z are as set forth in the following Table 1:






















Table 1

Compd.	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	Y	Z
A	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{CF}_3$	F
B	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{CF}_3$	
C	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{OCF}_3$	F
D	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{OCF}_3$	$-\text{NH}(\text{CH}_2)_2\text{OH}$











Compd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z
E	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	-CH <sub>3</sub>
F	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	F
G	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	H
H	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	F
I	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	H
J	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> -			H	F

Compd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z
K		-S(O <sub>2</sub> )-			Cl	F
L	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H
M	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-N(CH <sub>3</sub> ) <sub>2</sub>	F
N	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H
O	-N(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H
P	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-N(CH <sub>3</sub> ) <sub>2</sub>	H
Q	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H


Compd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z
R	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OH	F
S	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-				F
T	-CH(CH <sub>3</sub> ) <sub>2</sub>	Covalent bond			-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>3</sub>
U	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-				H
V		-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H
W		-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	H
X	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H

Compd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z
Y	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H
Z		-S(O <sub>2</sub> )-				H
AA		-S(O <sub>2</sub> )-				H
AB	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CN	H
AC	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-				H
AD	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CF <sub>3</sub>	H
AE	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	F
AF	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	F

Compd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z
AG	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	H
AH		-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	F
AI		-S(O <sub>2</sub> )-				H
AJ	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H
AK	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH(CH <sub>3</sub> ) <sub>2</sub>	F
AL	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH(CH <sub>3</sub> ) <sub>2</sub>	H
AM	-OCH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H
AN	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH(CH <sub>3</sub> ) <sub>2</sub>	-COOCH <sub>3</sub>

Compd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z
AO	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	F
AP	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H
AQ	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H
AR	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CF <sub>3</sub>	H
AS	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H

Additional preferred compounds are represented by formula III:



III

or a pharmaceutically acceptable salt thereof, wherein




Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ , cycloalkyl, halogen,  $-\text{OCF}_3$  and  $-\text{OH}$ ;

Z is selected from the group consisting of hydrogen, alkyl,  $-\text{CF}_3$ , halogen,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$  and  $-\text{OH}$ ;

5 and

$\text{M}^2$  is aryl or heteroaryl.

Additional preferred compounds are represented by formula IV:



IV

10

or a pharmaceutically acceptable salt thereof, wherein

Y is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-\text{OCF}_3$ ;

Z is selected from the group consisting of hydrogen, alkyl,  $-\text{CF}_3$ , halogen,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$  and  $-\text{OH}$ ;


15

and

$\text{M}^2$  is aryl or heteroaryl.

Additional preferred compounds are represented by formula V:

17



V

or a pharmaceutically acceptable salt thereof, wherein


Z is selected from the group consisting of hydrogen, -CF<sub>3</sub>, halogen,

5 -OCF<sub>3</sub> and -OH;

and

M<sup>2</sup> is aryl or heteroaryl.

Additional preferred compounds are represented by formula VI:



VI

10

or a pharmaceutically acceptable salt thereof, wherein





M<sup>2</sup> is aryl or heteroaryl;

and

Z is selected from the group consisting of hydrogen, alkyl, halogen, -CF<sub>3</sub>,

15 -N(R<sup>2</sup>)<sub>2</sub>, -OH and -OCF<sub>3</sub>.

Still additional preferred compounds are represented by the following formulae:



5 As used above and throughout this disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

"Patient" includes both human and animals.

"Mammal" means humans and other mammalian animals.

10 "Alkyl" means an aliphatic hydrocarbon group, which may be straight or branched and comprising about 1 to about 20 carbon atoms in the chain. Preferred alkyl groups contain about 1 to about 12 carbon atoms in the chain. More preferred alkyl groups contain about 1 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. "Lower alkyl" means a group having about 1 to about 6 carbon  
 15 atoms in the chain which may be straight or branched. The term "substituted alkyl" means that the alkyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the

group consisting of halo, alkyl, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy and -C(O)O-alkyl. Non-limiting examples of suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl and t-butyl.


"Alkynyl" means an aliphatic hydrocarbon group containing at least one carbon-  
5 carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain, and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkynyl chain. "Lower alkynyl" means about 2 to  
10 about 6 carbon atoms in the chain which may be straight or branched. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butynyl and 3-methylbutynyl. The term "substituted alkynyl" means that the alkynyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of alkyl, aryl and  
15 cycloalkyl.

"Alkylene" means a difunctional group obtained by removal of a hydrogen atom from an alkyl group that is defined above. Non-limiting examples of alkylene include methylene, ethylene and propylene.

"Aryl" means an aromatic monocyclic or multicyclic ring system comprising  
20 about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl.

"Heteroaryl" means an aromatic monocyclic or multicyclic ring system  
25 comprising about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the ring atoms is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. Preferred heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as  
30 defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. A nitrogen atom of a heteroaryl can be optionally oxidized to the corresponding N-oxide. Non-limiting examples of suitable heteroaryls include pyridyl, pyridyl-N-oxide,

pyrazinyl, furanyl, thiophenyl, pyrimidinyl, isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, 1,2,4-thiadiazolyl, pyrazinyl, pyridazinyl, quinoxaliny, phthalazinyl, imidazo[1,2-a]pyridinyl, imidazo[2,1-b]thiazolyl, benzofurazanyl, indolyl, azaindolyl, benzimidazolyl, benzothienyl, quinolinyl, 5 imidazolyl, thienopyridyl, quinazolinyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, isoquinolinyl, benzoazaindolyl, 1,2,4-triazinyl, benzothiazolyl and the like.




"Aralkyl" or "arylalkyl" means an aryl-alkyl- group in which the aryl and alkyl are 10 as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, 2-phenethyl and naphthalenylmethyl. The bond to the parent moiety is through the alkyl.

"Alkylaryl" means an alkyl-aryl- group in which the alkyl and aryl are as 15 previously described. Preferred alkylaryls comprise a lower alkyl group. Non-limiting example of a suitable alkylaryl group is tolyl. The bond to the parent moiety is through the aryl.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms. The cycloalkyl can be 20 optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkyls include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable multicyclic cycloalkyls include 1-decalinyl, norbornyl, adamantyl and the like. "Cycloalkenyl" means a non-aromatic mono- or 25 multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms, and which contains at least one carbon-carbon double bond. The cycloalkenyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above.

"Halogen" or "halo" means fluorine, chlorine, bromine, or iodine. Preferred are 30 fluorine, chlorine or bromine, and more preferred are fluorine and chlorine.

"Ring system substituent" means a substituent attached to an aromatic or non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being independently selected from the group consisting of aryl, heteroaryl, aralkyl, alkylaryl, aralkenyl, heteroaralkyl, alkylheteroaryl, heteroaralkenyl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halo, nitro, cyano, carboxy, alkoxycarbonyl, aryloxy carbonyl, aralkoxy carbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylsulfinyl, arylsulfinyl, heteroarylsulfinyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl,  $Y_1Y_2N-$ ,  $Y_1Y_2N$ -alkyl-,  $Y_1Y_2NC(O)-$  and  $Y_1Y_2NSO_2-$ , wherein  $Y_1$  and  $Y_2$  may be the same or different and are independently selected from the group consisting of hydrogen, alkyl, aryl, and aralkyl. "Ring system substituent" also means a cyclic ring of 3 to 7 ring atoms of which 1-2 may be a heteroatom, attached to an aryl, heteroaryl, heterocyclyl or heterocyclenyl ring by simultaneously substituting two ring hydrogen atoms on said aryl, heteroaryl, heterocyclyl or heterocyclenyl ring. Non-limiting examples include:



and the like.

"Heterocyclenyl" means a non-aromatic monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination, and which contains at least one carbon-carbon double bond or carbon-nitrogen double bond. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclenyl rings contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclenyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclenyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. The nitrogen or sulfur atom of the heterocyclenyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting

examples of suitable monocyclic azaheterocyclenyl groups include 1,2,3,4-tetrahydropyridine, 1,2-dihydropyridyl, 1,4-dihydropyridyl, 1,2,3,6-tetrahydropyridine, 1,4,5,6-tetrahydropyrimidine, 2-pyrrolinyl, 3-pyrrolinyl, 2-imidazoliny, 2-pyrazoliny, and the like. Non-limiting examples of suitable oxaheterocyclenyl groups include 3,4-dihydro-2H-pyran, dihydrofuranyl, fluorodihydrofuranyl, and the like. Non-limiting example of a suitable multicyclic oxaheterocyclenyl group is 7-oxabicyclo[2.2.1]heptenyl. Non-limiting examples of suitable monocyclic thiaheterocyclenyl rings include dihydrothiophenyl, dihydrothiopyranyl, and the like.

"Heterocyclyl" (or "heterocyclic") means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, and the like.

"Heteroaralkyl" means a heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain a lower alkyl group. Non-limiting examples of suitable aralkyl groups include pyridylmethyl, and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkyl.

"Hydroxyalkyl" means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

"Acyl" means an H-C(O)-, alkyl-C(O)- or cycloalkyl-C(O)-, group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl and propanoyl.

"Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1- naphthoyl.

5 "Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy, n-propoxy, isopropoxy and n-butoxy. The bond to the parent moiety is through the ether oxygen.

10 "Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond to the parent moiety is through the ether oxygen.

"Aralkyloxy" means an aralkyl-O- group in which the aralkyl group is as previously described. Non-limiting examples of suitable aralkyloxy groups include benzyloxy and 1- or 2-naphthalenemethoxy. The bond to the parent moiety is through the ether oxygen.

15 "Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkylthio groups include methylthio and ethylthio. The bond to the parent moiety is through the sulfur.

20 "Arylthio" means an aryl-S- group in which the aryl group is as previously described. Non-limiting examples of suitable arylthio groups include phenylthio and naphthylthio. The bond to the parent moiety is through the sulfur.

"Aralkylthio" means an aralkyl-S- group in which the aralkyl group is as previously described. Non-limiting example of a suitable aralkylthio group is benzylthio. The bond to the parent moiety is through the sulfur.

25 "Alkoxycarbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxycarbonyl groups include methoxycarbonyl and ethoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aryloxycarbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxycarbonyl groups include phenoxycarbonyl and naphthoxycarbonyl. The bond to the parent moiety is through the carbonyl.

30 "Aralkoxycarbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxycarbonyl group is benzyloxycarbonyl. The bond to the parent moiety is through the carbonyl.



"Alkylsulfonyl" means an alkyl-S(O<sub>2</sub>)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

5 "Arylsulfonyl" means an aryl-S(O<sub>2</sub>)- group. The bond to the parent moiety is through the sulfonyl.

"Halogenated alkyl" or "haloalkyl" means alkyl having 1 or more halogen atoms.

"Heteroalkyl" means straight or branched alkyl chain comprised of from 1 to 12 carbon atoms and 1 or more heteroatoms independently selected from the group consisting of N, O and S.

10 The term "substituted" means that one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

15 By "stable compound" or "stable structure" is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties.

20 It should also be noted that any heteroatom with unsatisfied valences in the text, schemes, examples and Tables herein is assumed to have a hydrogen atom to satisfy the valences.

When a functional group in a compound is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site

25 when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for example, T. W. Greene *et al*, *Protective Groups in Organic Synthesis* (1991), Wiley, New York, which is incorporated herein by reference thereto.

When any variable (e.g., aryl, heterocycle, R<sup>2</sup>, etc.) occurs more than one time

30 in any constituent or in Formulas I, II, III, IV, V or VI, its definition on each occurrence is independent of its definition at every other occurrence.

As a general note to all the Tables that are attached hereto as well as to the Description, Examples and Schemes in this application, any open-ended nitrogen

atom with unfulfilled valence in the chemical structures herein refers to NH, or in the case of a terminal nitrogen, -NH<sub>2</sub>. Similarly, any open-ended oxygen atom with unfulfilled valence in the chemical structures herein refers to -OH and any open-ended carbon atom with unfilled valence is appropriately filled with -H.

5           As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from the combination of the specified ingredients in the specified amounts.

          Prodrugs and solvates of the compounds of the invention are also  
10       contemplated herein. The term "prodrug", as employed herein, denotes a compound that is a drug precursor which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of Formulas I, II, III, IV, V or VI or a salt and/or solvate thereof. A discussion of prodrugs is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems* (1987) 14 of the A.C.S.  
15       Symposium Series, and in *Bioreversible Carriers in Drug Design*, (1987) Edward B. Roche, ed., American Pharmaceutical Association and Pergamon Press, both of which are incorporated herein by reference thereto.

          "Solvate" means a physical association of a compound of this invention with one or more solvent molecules. This physical association involves varying degrees of  
20       ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanولات, methanولات, and the like. "Hydrate" is a solvate  
25       wherein the solvent molecule is H<sub>2</sub>O.

          "Effective amount" or "therapeutically effective amount" is meant to describe an amount of compound or a composition of the present invention effective in inhibiting the CDK(s) and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect.

30       The compounds of Formulas I, II, III, IV, V or VI can form salts that are also within the scope of this invention. Reference to a compound of Formulas I, II, III, IV, V or VI herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with

inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of either Formulas I, II, III, IV, V or VI contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compounds of Formulas I, II, III, IV, V or VI may be formed, for example, by reacting a compound of either Formulas I, II, III, IV, V or VI with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides, lactates, maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates) and the like. Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by S. Berge *et al*, *Journal of Pharmaceutical Sciences* (1977) 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33 201-217; Anderson *et al*, *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as dicyclohexylamines, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, and dibutyl sulfates), long chain halides (e.g. decyl, lauryl, and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

All such acid salts and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.

5           Compounds of Formulas I, II, III, IV, V or VI, and salts, solvates and prodrugs thereof, may exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

          All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates and prodrugs of the  
10       compounds as well as the salts and solvates of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this invention, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl).  
15       Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the *IUPAC 1974* Recommendations. The use of the terms "salt", "solvate" "prodrug" and the like, is  
20       intended to equally apply to the salt, solvate and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the inventive compounds.

          The compounds of the present invention can be useful as cannabinoid receptor ligands. The compounds can have anti-inflammatory activity and/or  
25       immunomodulatory activity and can be useful in the treatment of various medical conditions including, e.g., cutaneous T-cell lymphoma, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, sepsis, shock, sarcoidosis, idiopathic pulmonary fibrosis, bronchopulmonary dysplasia, retinal disease, scleroderma, osteoporosis, renal ischemia, myocardial infarction, cerebral  
30       stroke, cerebral ischemia, nephritis, hepatitis, glomerulonephritis, cryptogenic fibrosing aveolitis, psoriasis, transplant rejection, atopic dermatitis, vasculitis, allergy, seasonal allergic rhinitis, Crohn's disease, inflammatory bowel disease, reversible airway obstruction, adult respiratory distress syndrome, asthma, chronic obstructive

pulmonary disease (COPD) or bronchitis. It is contemplated that a compound of this invention may be useful in treating one or more of the diseases listed.

Additionally, a compound of the present invention may be co-administered or used in combination with disease-modifying antirheumatic drugs (DMARDS) such as methotrexate, azathioprine, leflunomide, pencillinamine, gold salts, mycophenolate mofetil, cyclophosphamide and other similar drugs. They may also be co-administered with or used in combination with NSAIDS such as piroxicam, naproxen, indomethacin, ibuprofen and the like; COX-2 selective inhibitors such as Vioxx® and Celebrex®; immunosuppressives such as steroids, cyclosporin, Tacrolimus, rapamycin and the like; biological response modifiers (BRMs) such as Enbrel®, Remicade®, IL-1 antagonists, anti-CD40, anti-CD28, IL-10, anti-adhesion molecules and the like; and other anti-inflammatory agents such as p38 kinase inhibitors, PDE4 inhibitors, TACE inhibitors, chemokine receptor antagonists, Thalidomide and other small molecule inhibitors of pro-inflammatory cytokine production.

Also additionally, one or more compounds of the present invention may be co-administered or used in combination with at least one H1 antagonist for the treatment of seasonal allergic rhinitis and/or asthma. Suitable H1 antagonists may be, for example, Claritin®, Clarinex®, Allegra®, or Zyrtec®.

In another aspect, the invention provides a method for treating rheumatoid arthritis comprising administering at least one compound of either Formulas I, II, III, IV, V or VI in combination with at least one compound selected from the class consisting of a COX-2 inhibitor e.g. Celebrex® or Vioxx®; a COX-1 inhibitor e.g. Feldene®; an immunosuppressive e.g. methotrexate or cyclosporin; a steroid e.g.  $\beta$ -methasone; and anti-TNF- $\alpha$  compound, e.g. Enbrel® or Remicade®; a PDE IV inhibitor, or other classes of compounds indicated for the treatment of rheumatoid arthritis.

In another aspect, the invention provides a method for treating multiple sclerosis comprising administering at least one compound of either Formulas I, II, III, IV, V or VI in combination with at least one compound selected from the group consisting of Avonex®, Betaseron, Copaxone or other compounds indicated for the treatment of multiple sclerosis.

In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of at least one compound of Formula I, II, III, IV, V or VI in at least one pharmaceutically acceptable carrier.

5 In yet another aspect, the invention relates to a pharmaceutical composition for treating rheumatoid arthritis comprising a therapeutically effective amount of at least one compound of Formula I, II, III, IV, V or VI in combination with at least one compound selected from the class consisting of a COX-2 inhibitor, a COX-1 inhibitor, an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of rheumatoid arthritis.

10 In still another aspect, the invention relates to a pharmaceutical composition for treating multiple sclerosis comprising a therapeutically effective amount of at least one compound of Formula I, II, III, IV, V or VI in combination with at least one compound selected from the group consisting of Avonex®, Betaseron, Copaxone or other compounds indicated for the treatment of multiple sclerosis.

15 The compounds of the present invention exhibit anti-inflammatory and/or immunomodulatory activity and can be useful in the treatment of various medical conditions including, e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, osteoporosis, renal ischemia, cerebral stroke, cerebral ischemia, nephritis, psoriasis, allergy, inflammatory disorders of the lungs and  
20 gastrointestinal tract such as Crohn's disease, and respiratory tract disorders such as reversible airway obstruction, asthma, chronic obstructive pulmonary disease (COPD) and bronchitis. This utility is manifested as demonstrated by activity in the following assay.

Potential cannabinoid receptor ligands were screened for the ability to compete  
25 with [ $^3$ H] CP-55,940 for binding to recombinant cannabinoid receptors. Test compounds were serially diluted in Diluent Buffer (50 mM Tris pH 7.1, 1 mM EDTA, 3 mM MgCl<sub>2</sub>, 0.1% BSA, 10% DMSO, 0.36% methyl cellulose (Sigma M-6385)) from stocks prepared in 100% DMSO. Aliquots (10  $\mu$ L) were transferred into 96-well microtiter plates. Membrane preparations of recombinant human cannabinoid CB<sub>2</sub>  
30 receptor (Receptor Biology #RB-HCB2) or recombinant human cannabinoid CB<sub>1</sub> receptor (Receptor Biology #RB-HCB1) were diluted to 0.3 mg/mL in Binding Buffer (50 mM Tris pH 7.2, 1 mM EDTA, 3 mM MgCl<sub>2</sub>, 0.1% BSA). Aliquots (50  $\mu$ L) were

added to each well of the microtiter plate. The binding reactions were initiated by addition of [ $^3\text{H}$ ] CP-55,940 (New England Nuclear # NET 1051; specific activity =180 Ci/mmol) to each well of the microtiter plate. Each 100  $\mu\text{l}$  reaction mixture contained 0.48 nM [ $^3\text{H}$ ] CP-55,940, 15  $\mu\text{g}$  membrane protein in binding buffer containing 1%  
5 DMSO and 0.036 % methyl cellulose. Following incubation for 2 hours at room temperature, the reactions were filtered through 0.5% polyethylenimine-coated GF/C filter plates (UniFilter-96, Packard) with a TomTec Mark 3U Harvester (Hamden, CT). The filter plate was washed 5 times with binding buffer, rotated 180°, then re-washed 5 times with binding buffer. Bound radioactivity was quantitated following addition of  
10 30  $\mu\text{l}$  of Packard Microscint 20 scintillant in a Packard TopCount NXT microplate scintillation counter. Non-linear regression analysis of the resulting data was performed using Prism 2.0b (GraphPad, San Diego, CA).

The compounds of the invention exhibit potent affinities for the CB<sub>2</sub> receptor as measured by K<sub>i</sub> values (in nM). The activities (potencies) for the compounds of the  
15 invention are determined by measuring their K<sub>i</sub> values. The smaller the K<sub>i</sub> value, the more active the compound is for modulating the CB<sub>2</sub> receptor. Compounds of the invention exhibit a wide range of activities. The CB<sub>2</sub> average K<sub>i</sub> values for compounds having one of the Formulae I, II, III, IV, V or VI generally range from > 0 nM (e.g., 0.1 nM) to about 1000 nM, preferably about 0.1 nM to about 1000 nM, more preferably  
20 about 0.1 nM to about 100 nM, more preferably about 0.1 to about 20 nM, and most preferably less than about 20 nM. Representative compounds of the invention that exhibit excellent CB<sub>2</sub> inhibitory activity (K<sub>i</sub> values of less than about 20 nanomolar, nM) are as follows: Compounds C, F, G, K, S, AG, AH, AI, AJ, AK, AL, AO, and AQ from Table 1.

25 The inventive compounds are also highly selective for modulating a CB<sub>2</sub> receptor as opposed to modulating a CB<sub>1</sub> receptor. A "selective modulator" means that a compound's selection ratio of K<sub>i</sub> of the CB<sub>1</sub> receptor to the K<sub>i</sub> of the CB<sub>2</sub> receptor is greater than about 100, preferably greater than about 500, more preferably greater than about 1000 and most preferably greater than about 3000.

30 Compounds of the present invention were found to exhibit CB<sub>2</sub> receptor binding activity in the range of 0.1 to 1000 nM.

The present invention also relates to a pharmaceutical composition comprising one or more compounds of Formula I, II, III, IV, V or VI of this invention and one or

more pharmaceutically acceptable carriers. The compound of Formulas I, II, III, IV, V or VI, can be administered in any conventional dosage form known to those skilled in the art. Pharmaceutical compositions containing the compound of Formula I, II, III, IV, V or VI can be prepared using conventional pharmaceutically acceptable excipients and additives and conventional techniques. Such pharmaceutically acceptable excipients and additives include non-toxic compatible fillers, binders, disintegrants, buffers, preservatives, anti-oxidants, lubricants, flavorings, thickeners, coloring agents, emulsifiers and the like. All routes of administration are contemplated including, but not limited to, parenteral, transdermal, subcutaneous, intramuscular, sublingual, inhalation, rectal and topical.

Thus, appropriate unit forms of administration include oral forms such as tablets, capsules, powders, cachets, granules and solutions or suspensions, sublingual and buccal forms of administration, aerosols, implants, subcutaneous, intramuscular, intravenous, intranasal, intraocular, subcutaneous or rectal forms of administration.

When a solid composition is prepared in the form of tablets, e.g., a wetting agent such as sodium lauryl sulfate can be added to micronized or non-micronized compound of Formula I, II, III, IV, V or VI and mixed with a pharmaceutical vehicle such as silica, gelatin starch, lactose, magnesium stearate, talc, gum arabic or the like. The tablets can be coated with sucrose, various polymers or other appropriate substances. Tablets can be treated so as to have a prolonged or delayed activity and so as to release a predetermined amount of active principle continuously or at predetermined intervals, e.g., by using ionic resins and the like.

A preparation in the form of gelatin capsules may be obtained, e.g., by mixing the active principle with a diluent, such as a glycol or a glycerol ester, and incorporating the resulting mixture into soft or hard gelatin capsules.

A preparation in the form of a syrup or elixir can contain the active principle together, e.g., with a sweetener, methylparaben and propylparaben as antiseptics, flavoring agents and an appropriate color.

Water-dispersible powders or granules can contain the active principle mixed, e.g., with dispersants, wetting agents or suspending agents, such as polyvinylpyrrolidone, as well as with sweeteners and/or other flavoring agents.



Rectal administration may be provided by using suppositories which may be prepared, e.g., with binders melting at the rectal temperature, for example cocoa butter or polyethylene glycols.

5 Parenteral, intranasal or intraocular administration may be provided by using, e.g., aqueous suspensions, isotonic saline solutions or sterile and injectable solutions containing pharmacologically compatible dispersants and/or solubilizers, for example, propylene glycol or polyethylene glycol.

10 Thus, to prepare an aqueous solution for intravenous injection, it is possible to use a co-solvent, e.g., an alcohol such as ethanol or a glycol such as polyethylene glycol or propylene glycol, and a hydrophilic surfactant such as Tween® 80. An oily solution injectable intramuscularly can be prepared, e.g., by solubilizing the active principle with a triglyceride or a glycerol ester.

Topical administration can be provided by using, e.g., creams, ointments or gels.

15 Transdermal administration can be provided by using patches in the form of a multilaminate, or with a reservoir, containing the active principle and an appropriate solvent.

Administration by inhalation can be provided by using, e.g., an aerosol containing sorbitan trioleate or oleic acid, for example, together with  
20 trichlorofluoromethane, dichlorofluoromethane, dichlorotetrafluoroethane or any other biologically compatible propellant gas; it is also possible to use a system containing the active principle, by itself or associated with an excipient, in powder form.

The active principle can also be formulated as microcapsules or microspheres, e.g., liposomes, optionally with one or more carriers or additives.

25 Implants are among the prolonged release forms which can be used in the case of chronic treatments. They can be prepared in the form of an oily suspension or in the form of a suspension of microspheres in an isotonic medium.

30 The daily dose of a compound of Formula I, II, III, IV, V or VI for treatment of a disease or condition cited above is about 0.001 to about 100 mg/kg of body weight per day, preferably about 0.001 to about 10 mg/kg. For an average body weight of 70 kg, the dosage level is therefore from about 0.1 to about 700 mg of drug per day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the

attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient.

Compounds of the present invention can be used in combination with disease modifying antirheumatic agents described herein above, the administration and dosage of such agents is as according to the schedule listed in the product information sheet of the approved agents, in the Physicians Desk Reference (PDR) as well as therapeutic protocols well known in the art.

Compounds of the present invention can be used in combination with H1 antagonists described herein above, the administration and dosage of such agents is as according to the schedule listed in the product information sheet of the approved agents, in the Physicians Desk Reference (PDR) as well as therapeutic protocols well known in the art.

Compounds of the present invention can be used in combination with compounds useful in the treatment of multiple sclerosis described herein above, the administration and dosage of such agents is as according to the schedule listed in the product information sheet of the approved agents, in the *Physicians Desk Reference* (PDR) as well as therapeutic protocols well known in the art.

Compounds of the present invention can be used in combination with compounds useful in the treatment of psoriasis described herein above, the administration and dosage of such agents is as according to the schedule listed in the product information sheet of the approved agents, in the Physicians Desk Reference (PDR) as well as therapeutic protocols well known in the art.

It will be apparent to those skilled in the art that the administration of the agents used in combination with the compounds of the present invention can be varied depending on the disease being treated and the known effects of the agents on that disease. Also, in accordance with the knowledge of the skilled clinician, the therapeutic protocols (e.g. dosage amounts and times of administration) can be varied in view of the observed effects of the administered agents on the patients, and in view of the observed responses of the disease to the administered agents.

Compounds of the present invention are generally prepared by processes known in the art, for example, by the processes described below.

The following abbreviations are used in the procedures and schemes: aqueous (aq.), anhydrous (anhyd), *n*-butyl (*n*-Bu), *n*-butyllithium (*n*-BuLi), tert-butyllithium (t-


BuLi), butyllithium (BuLi), sodium t-butoxide ( $\text{NaOBu}^t$ ), concentrated (conc.), diethyl ether ( $\text{Et}_2\text{O}$ ), days (d), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), dimethylformamide (DMF), ethanol (EtOH), ethyl (Et), ethyl acetate (EtOAc), hours (h), leaving group (LG), hydroxybenzotriazole (HOBT), *meta*-chloroperoxybenzoic acid (mCPBA), lithium diisopropylamide (LDA), methanesulfonyl chloride (MsCl), methanol (MeOH), palladium bis(dibenzylideneacetone) ( $\text{Pd}(\text{dba})_2$ ), minutes (min), methyl (Me), methyllithium (MeLi), molar (moles per liter, M), *N*-chlorosuccinimide (NCS), *N,N*-dimethylaminopyridine (DMAP), normal (N), pounds per square inch (psi), preparative thin layer chromatography (PTLC), thin layer chromatography (TLC), room temperature (rt), saturated sodium chloride solution (brine), milliliter (mL), millimole (i.e., 0.001 mole; mmol), milligram (mg), silica gel chromatography (sgc), 2-(*tert*-butoxycarbonyloxyimino)-2-phenylacetonitrile (BOC-ON), *tert*-butoxycarbonyl (BOC), trifluoroacetic anhydride (TFAA), trifluoroacetic acid (TFA), trifluoromethanesulfonic anhydride ( $\text{Tf}_2\text{O}$ ), and tetrahydrofuran (THF). In a typical work-up procedure, the reaction mixture is diluted with a suitable solvent, such as EtOAc,  $\text{Et}_2\text{O}$ , or  $\text{CH}_2\text{Cl}_2$ , and washed successively with appropriate acidic, basic, or neutral aqueous solutions. The organic solution is separated, dried over an appropriate drying agent such as  $\text{MgSO}_4$  or  $\text{Na}_2\text{SO}_4$ , filtered, and the solvent removed by evaporation.

### General Description of Methods

Compounds of formula I may be preferably prepared as shown in the following reaction schemes. They are preferably prepared as shown in the following reaction schemes. In the reaction schemes, the variables are defined as above.

## General Scheme 1

For compounds wherein  $L^1 = -CH_2-$ ,  $-C(O)-$ ,  $-C=NOR^2-$  and  $L^2 = -S(O_2)-$ ,  $-CH_2-$ ,  $-C(O)-$




Compounds wherein  $L^1$  is one of the following:  $-CH_2-$ ,  $-C(O)-$ ,  $-C=NOR^2-$  and  $L^2$  is one of the following:  $-S(O_2)-$ ,  $-CH_2-$ ,  $-C(O)-$  can be made as shown in Scheme 1.

## General Scheme 1

- 5 In step 1, an aldehyde of the formula 1 is treated with a base such as LDA, in the presence of N,N,N' trimethylethylenediamine in an inert organic solvent such as THF or DME to generate an ortho anion, and an electrophile of the formula 2 where FG is one of the functional groups e.g., sulfonylfluoride, disulfide, acid chloride, alkylbromide or aldehyde. Preferably, the reaction temperature ranges from  $-70^\circ\text{C}$  to  $0^\circ\text{C}$ . The product is purified by silica gel chromatography. In step 2, an aryl bromide of the formula 4 is subjected to halogen metal exchange with a base such as n-BuLi or t-BuLi in an inert solvent such as THF or ether at a low temperature in the range of  $-78^\circ\text{C}$  to  $-40^\circ\text{C}$  and treated with aldehyde 3 to give an intermediate compound of formula 5. In step 3, intermediate 5 is reduced with triethylsilane in the presence of trifluoroacetic acid to provide a target compound of the formula 6. Additionally, intermediate 5 can be oxidized, e.g., with PCC in methylene chloride, to provide another target compound of the general formula 7. Treatment of 7 with an O-alkylated hydroxyl amine in (excess) pyridine provides oxime product 8.

## General Scheme 2

For compounds wherein  $L^1 = -S(O_2)-$  and  $L^2 = -CH_2-, -C(O)-, -S(O_2)-$ 


Compounds wherein  $L^1$  is  $-S(O_2)-$  and  $L^2$  is one of the following:  $-S(O_2)-$ ,  $-CH_2-$ , or  $-C(O)-$  can be prepared as shown in Scheme 2.

## General Scheme 2

- 5 In step 1, aryl halide 9 is treated with an alkyl lithium such as n-butyllithium or t-butyllithium in an inert organic solvent such as THF or DME at  $-78^{\circ}\text{C}$ . The resulting aryllithium is then treated with sulfonyl fluoride (i) to provide sulfone product 11. Alternatively, the aryllithium is treated with a disulfide (ii) as shown in step 1a to provide sulfide 10, which is then oxidized with a reagent such as mCPBA (step 1 b) to provide product 11. In step 2, product 11 is treated with an alkyl lithium such as n-BuLi or t-BuLi and the resulting ortho anion of 11 is quenched with sulfonyl fluoride (iii) to provide the target molecule 12. Compound 12 can also be prepared by treating the ortho anion of 11 with a disulfide (iv) as shown in step 2a followed by oxidation of the resulting sulfide 13 with mCPBA. Additionally, the ortho anion can be quenched with an aldehyde to obtain alcohol 16 (step 3). In step 4, alcohol 16 can be oxidized to the target 14. The reduced form of alcohol 16, compound 15, can be obtained by using a
- 10
- 15

reducing agent such as triethylsilane in the presence of a strong acid such as trifluoroacetic as shown in step 5.


Compounds wherein  $L^1$  is  $-S(O_2)-$  and  $L^2$  is  $-S(O_2)-$  are prepared as shown in Scheme 3.



### General Scheme 3

In step 1, starting material 17 is treated with a base such as BuLi, NaH, or CsCO<sub>3</sub> followed by compound 18, where LG is a leaving group, preferably a fluoride. The product 19 is treated with an alkyl lithium such as nBuLi or t-BuLi at a low temperature, e.g.,  $-100^{\circ}\text{C}$  to  $-78^{\circ}\text{C}$ , in an inert solvent such as THF or DME, and quenched with sulfonyl fluoride 20 to provide target compound 21. Alternatively, a disulfide 24 is used to quench the anion, thereby providing compound 25. Compound 25 is then oxidized with an oxidizing agent, e.g., mCPBA, to the target 21. Another route to access target 21 involves quenching the anion of 19 with SO<sub>2</sub> and treating the resulting lithium salt of sulfonic acid with chlorinating agent such as N-chlorosuccinimide, thereby forming a sulfonyl chloride, which is converted to the corresponding fluoride 22. Compound 22 is then reacted with an aryl lithium reagent formed by reacting 23 with alkyl lithium at  $-78^{\circ}\text{C}$  to give target 21.

38




#### Preparation of Dihydrofuran compounds: General Scheme 4

In **step 1**, 2,4-dibromophenol **26** and dibromoethane are dissolved in a suitable solvent, such as methylene chloride, dichloroethane, or toluene. Aqueous NaOH is used as a base. The reaction mixture is stirred in the presence of a phase-transfer catalyst (such as tetrabutylammonium hydrogen sulfate, methyl-n-butylammonium chloride, or benzyltriethylammonium hydroxide) between 0 °C and 100 °C for 0.5 to 48 h. The product may be purified via silica gel chromatography or crystallization.

In **step 2**, the product of step 1 (**27**) is dissolved in THF or ether and treated with a base such as n-BuLi at -100 °C. After stirring, the reaction mixture is treated with another portion of n-BuLi at -78 °C. The resulting anion is trapped with the sulfonyl fluoride compound **28** prepared in General Scheme 1. The product can be purified via chromatography or crystallization.

In **step 3**, the product of step 2 (**29**) is dissolved in THF and treated with a base such as n-BuLi at -78 °C to form a dianion, which is trapped with a suitable electrophile (such as **iii** / **iv**). The reaction mixture is quenched with a suitable proton source such as aq NH<sub>4</sub>Cl or phosphate buffer then extracted with EtOAc. The product **30** may be purified via silica gel chromatography or crystallization.

## General Scheme 5



## General Scheme 5

In **step 1**, a mixture of 1,1,2 trichloroethane **31**, substituted thiophenol **32**, 1 N NaOH and a phase transfer catalyst (such as Aliquat 336, tetrabutylammonium hydrogen sulfate) is stirred for four days. The mixture is then treated with an oxidizing agent (such as  $\text{H}_2\text{O}_2$  or mCPBA) and a catalytic amount of ammonium molybdate. The reaction is then diluted with an organic solvent such as ethyl acetate, methylene chloride, or ether. The organic phase is washed with water and brine and dried. The product **33** is isolated by silica-gel chromatography.


In **step 2**, product **33** dissolved in methylene chloride at a temperature of  $0^\circ\text{C}$ - $15^\circ\text{C}$  is treated with a base such as triethyl amine, diisopropylamine, or DBU. After stirring for 1 h, the reaction is washed with water, and brine. The product **34** is isolated with silica-gel chromatography.

In **step 3**, the thiol **35** is dissolved in a solvent such as THF or ether, cooled to  $0^\circ\text{C}$  and treated with a base such as sodium hydride. After stirring for 10-30 min, compound **34** is added as a THF or ether solution and the resulting mixture is allowed to warm to room temperature and the reaction is stirred for 10h. The reaction is quenched with water and worked up. The product is isolated by silica-gel chromatography to provide product **36**.

In **step 4**, product **36** is dissolved in glacial acetic acid and treated with an oxidizing agent such as hydrogen peroxide at  $40$ - $60^\circ\text{C}$  and stirred for 3h. The reaction is cooled and diluted with an organic solvent such as ethylacetate, methylene chloride, or ether and worked up. The product **37** is recrystallized from ether as a white solid.



### Synthesis of Compound 6B



**Step 1)** n-BuLi (0.66 mL, 2.5 M in hexanes) was added to a solution of 4-bromoisopropylbenzene **4A** (0.3 g, 16 mmol) in anhydrous THF (3 mL) at  $-78^{\circ}\text{C}$ . The mixture was stirred for 20 minutes and then a THF solution of aldehyde **3A** (0.5 g, 1.6 mmol) was added while maintaining the temperature at  $-78^{\circ}\text{C}$ . The resulting mixture was stirred for 2h. The reaction was quenched with water, and allowed to warm to  $0^{\circ}\text{C}$ . The product was extracted with 50 mL ethylacetate. The organic phases were washed with 2 X 25 mL water and 1 x 25 mL brine and dried over  $\text{Na}_2\text{SO}_4$ . The

solvent was removed, and the crude product was subjected silica gel preparatory plate chromatography using 20% ethyl acetate / hexane as the eluting solvent to provide 0.5 g of pure product **5A** (alcohol).


**Step 1A)** Aldehyde **3A** was prepared using the following procedure: To a solution of N,N,N'-Trimethylethylenediamine (1.2 mL, 8.6 mmols) in THF (8 mL) at -20°C was added n-BuLi (1.6 M, 5.4 mL, 8.6 mmol) dropwise. After 15 minutes 4-trifluoromethoxybenzaldehyde (**1A**) (1.5 g, 7.8 mmol) in THF (8 mL) was added. The mixture was stirred for 15 minutes and more n-BuLi (1.6M, 14.6 mL, 23 mmol) was added. The reaction mixture was stirred at -20°C for 1h, and then placed in the freezer at -20°C for 20h. The next day, the mixture was cooled to -40°C, followed by addition of a solution of o-fluorobenzenedisulfide (**2A**) (4.0 g, 15.7 mmol) in 30 mL THF and stirred at -40°C to -35°C for 3 h. The reaction was poured into 0.5 N HCl and extracted with ethyl acetate. The organic layer was washed with water and brine and then dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give a yellow oil. The oil was purified by silica-gel column chromatography (3 % ethylacetate / hexanes) to give a light yellow solid **3A**, yield 1.55g (62 %).

**Step 2)** The product **5A** from **step 1** (0.5 g, 0.001 mol) in 50 mL methylene chloride was treated with triethyl silane (1.83 mL, 0.01 mol) followed by dropwise addition of trifluoroacetic acid (0.097 mL, 0.0011 mol) at ambient temperature. The mixture was stirred for 10 h. The solvent and excess reagents were removed under reduced pressure. The resulting crude product was subjected to preparatory plate silica gel chromatography using ethylacetate : hexane (1:10) as the developing solvent, to provide 0.43 g of pure product **6A**.

**Step 3)** The product **6A** from **step 2** (0.4 g, 0.00095 mol) in 20 mL methylene chloride was treated with metachloroperoxy benzoic acid (0.87g, 77%, 0.0035 mol). The mixture was stirred for 20 h, then washed in sequence with 5% aq. sodium bisulfite (20 mL), aq. sodium bicarbonate (2 x 20 mL), and brine (2X 20 mL). The organic phases were dried and concentrated under reduced pressure to yield a crude product which was then purified by preparatory plate silica gel chromatography using ethylacetate : hexane (1:6) as the developing solvent, to provide 0.41 g of pure product **6B**.

**Synthesis of Compounds 6D and 6E :**

42




**Step 1:** Compound **6B** (0.3 g; 0.00067) in 20 mL acetonitrile was treated with N-Boc piperazine (0.6 g; 0.0033 mol) and refluxed for 24h. Additional N-Boc piperazine was added (0.6 g; 0.0033 mol) and refluxing was continued for another 24h. The reaction was cooled and the solvent was removed under reduced pressure. The product was isolated by preparatory silica-gel chromatography using ethylacetate:hexane (1:1) to obtain 0.32g of pure product **6C**.

**Step 2:** Product **6C** from **Step 1** (0.129g, 0.0002 mol) in 20 mL methylene chloride was treated with trifluoroacetic acid (0.05 mL, 0.0006 mol) and the mixture was stirred for 3 h at ambient temperature. The reaction was diluted with 20 mL methylene chloride and washed with 5 mL aq. sodium bicarbonate. The organic phases were dried and concentrated to yield pure product **6D** (0.08g).

**Synthesis of 6E:** Product **6B** (0.07 g, 0.000155 mol) dissolved in 10 mL acetonitrile was treated with ethanolamine (0.4 mL) and heated to 80°C and stirred for 24 h. The solvent was removed under reduced pressure and the crude product was re-dissolved in 30 ml methylene chloride and washed with water (1x 20 mL) and brine (1 x 20 mL). The organics were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The resulting crude was purified by preparatory plate silica-gel using ethylacetate : hexane (1:2) to give 0.04 g of pure product **6E**.

# Synthesis of Compounds 12A, 12B and 12C :



**Step1:** n-Butyl lithium (2.64 mL, 2.24M hexane solution, 0.006mol) was added dropwise at  $-70^{\circ}\text{C}$  to a dry THF solution (10 mL) of bromo isopropylbenzene **4A** (1g, 0.005 mol). The mixture was stirred for 0.5 h and then a solution of trifluoromethoxybenzene sulfonyl fluoride (1.34 g, 0.0055mol) as a THF solution (10mL) was added while maintaining the temperature at  $-70^{\circ}\text{C}$ . Stirring was continued for an additional 2 hours. The reaction was quenched with water (5 mL) and extracted with ethylacetate (100 mL). The organic phases were washed with water (2 x 50 mL) and brine (1 x 50 mL), dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to yield a crude product which was isolated by silica gel column chromatography using 10 % ethylacetate / hexane as the eluting solvent, resulting in 0.45g of the product **11A**.

**Step 2a:** The product **11A** (0.1 g, 0.00029 mol) of **step 1** was dissolved in anhydrous THF (10 mL) and cooled to  $-70^{\circ}\text{C}$  and treated with n-BuLi (0.116 mL, 2.5 M in hexanes, 0.00029 mol). The mixture was stirred for 0.5 h and treated with a THF solution of fluorophenyl disulfide (0.08 g, 0.003 mol, 5 mL THF). The resulting mixture was stirred for 2.5 h and the reaction was allowed to warm to  $-50^{\circ}\text{C}$  and then quenched with water (5 ml). The reaction mixture was warmed to room temperature and extracted with 50 mL ethylacetate. The organic phases were washed with water

(50 mL), brine (50 mL), and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solvent was removed under pressure and the crude product was subjected to silica-gel preparatory chromatography to provide 0.11 g of pure product **13A**.

**Step 3a:** The product **13A** from **step 2a** (0.066g, 0.00014 mol) was dissolved in  
5 methylene chloride (10 mL) and treated with metachloroperoxybenzoic acid (0.12 g, 77%, 0.0007 mol) as a methylene chloride (2 mL) solution. The mixture was stirred for 4 days. The reaction was washed with aq.  $\text{NaHCO}_3$  (2 x 20 mL), water (1 x 20 mL), and brine (1 x 20 mL). The organic phases were dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated to yield a crude product. The crude product was purified by silica gel  
10 preparatory chromatography using 10 % ethylacetate/hexane as the developing solvent to provide 0.039 g of pure product **12C**.

**Step 2b:** The product **11A** (0.1 g, 0.00029 mol) of **step 1** was dissolved in anhydrous THF (10 mL) and cooled to  $-70^\circ\text{C}$  and treated with n-BuLi (0.07 mL, 2.5 M in hexanes, 0.00016 mol). The mixture was stirred for 0.5 h and treated with a THF  
15 solution of pyridyl disulfide (0.035g, 0.00016 mol, 5 mL THF). The resulting mixture was stirred for 2 h and the reaction was allowed to warm to  $0^\circ\text{C}$  and then quenched with water (5 mL). The reaction mixture was warmed to room temperature and extracted with 50 mL ethylacetate. The organic phases were washed with water (50 mL), brine (50 mL), and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solvent was removed  
20 under reduced pressure and the crude product was subjected to silica-gel preparatory chromatography using 1:4 ethylacetate:hexane as the developing solvent to provide 0.073 g of pure product.


**Step 3b:** The product from **step 2b** (0.035g, 0.000077 mol) was dissolved in methylene chloride (10 mL) and treated with metachloroperoxybenzoic acid (0.076 g,  
25 77%, 0.0003 mol) as a methylene chloride (2 mL) solution. The mixture was stirred for 2 days. The reaction was washed with aq.  $\text{NaHCO}_3$  (2 x 20 mL), water (1 x 20 mL), and brine (1 x 20 mL). The organic phases were dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated to yield a crude product. The crude product was purified by silica gel preparatory chromatography using 10 % ethylacetate/hexane as the developing  
30 solvent to provide 0.006 g of pure product **12B** along with 0.01 g of over-oxidized product, the pyridine N-oxide.

**Step 2c:** The product **11A** of **step 1** (0.083 g, 0.00024 mol) was dissolved in anhydrous THF (10 mL) and cooled to  $-70^\circ\text{C}$  and treated with n-BuLi (0.106 mL, 2.5 M

in hexanes, 0.00026 mol). The mixture was stirred for 0.5 h and treated with a THF solution of methylfuranyldisulfide (0.065 g, 0.0028 mol, 5 mL THF). The resulting mixture was stirred for 2.5 h and the reaction was allowed to warm to -20°C and then quenched with water (5 mL). The reaction mixture was warmed to room temperature and extracted with 50 mL ethylacetate. The organic phases were washed with water (50 mL), brine (50 mL), and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solvent was removed under pressure and the crude product was subjected to silica-gel preparatory chromatography using 10% ethylacetate / hexane as the solvent to provide 0.09g of pure product.

**Step 3c:** The product from **step 2c** (0.09 g) was dissolved in methylene chloride (10 mL) and treated with metachloroperoxybenzoic acid (0.27 g) as a methylene chloride (2 mL) solution. The mixture was stirred for 4 days. The reaction was washed with aq. NaHCO<sub>3</sub> (2 x 20 mL), water (1 x 20 mL), and brine (1 x 20 mL). The organic phases were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a crude product. The crude product was purified by silica gel preparatory chromatography using 25 % ethylacetate/hexane as the developing solvent to provide 0.017 g of pure product **12A**.

### Synthesis of Compound 12 F



20


**Step 1:** Product **12C** from **step 3a** of the above scheme, (0.4g, 0.00079 mol) was dissolved in 50 mL of CH<sub>3</sub>OH and to this solution was added 1M aq. NaOH (1.8 mL). The resulting mixture was heated at 60°C for 16h. TLC analysis showed

incomplete reaction. Additional 1M aq. NaOH (0.9 mL) was added and refluxing was continued for 6h. The reaction was cooled and the solvent was removed under reduced pressure. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and washed with water (1X 50 mL) and brine (1 x 50 mL). The organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>),  
5 filtered, and concentrated to provide a crude product. The product was isolated by silica-gel preparatory plate chromatography using 30% ethylacetate in hexane to yield 0.3 g of pure product **12D**.

**Step 2:** Product **12D** from **step 1** (0.16g, 0.00035 mol) dissolved in methylene chloride (10 mL) was cooled to -10°C and treated dropwise with a 1 M solution of  
10 boron tribromide (1.32 mL) The mixture was allowed to warm up to room temperature and stirred for 16h. TLC analysis showed incomplete reaction. Additional boron tribromide (1.32 mL) was added and stirring was continued for 16 h. The reaction was quenched with 5 mL water and diluted with 50 mL methylene chloride. The organic phases were washed with water (1x 50 mL), aq. NaHCO<sub>3</sub> (1 x 50 mL), and brine (1 x  
15 50 mL). The organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to yield a crude product. The product **12E** was isolated by silica-gel preparatory plate chromatography using 30% ethylacetate/hexanes. Yield: 0.11g.

**Step 3:** Product **12E** from **Step 2** (0.039 g, 0.0000898 mol) in DMF (5 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (0.015g, 0.00011mol) followed by methoxybromoethane (0.0085  
20 mL, 0.000094 mol). The mixture was heated at 50°C for 24 h. The reaction was diluted with ethylacetate (50 mL) and the organic phases were washed with water (1x 50 mL) and saline (1x 50 mL). The organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to provide a crude product (compound **12F**), which was purified by preparatory plate chromatography using 30% ethylacetate /hexanes as the solvent.  
25 Yield: 0.009 g.

## Synthesis of Compounds 12G and 12H



**Step 1:** Dibromobenzene (5g, 0.021mol) was dissolved in 100 mL anhydrous THF and cooled to -70°C. *n*-BuLi (6.8 mL, 0.0168 mol, 2.5 M) was added and stirred for 1h, then a THF (10mL) solution of isopropylbenzene sulfonyl fluoride (4.3 g, 0.021 mol) was added and stirred for 3h, and the temperature was allowed to rise to -30°C. The reaction was quenched with water and the reaction was diluted with ethylacetate (100 mL). The organic phases were washed with water (1 x 50 mL) and brine (1 X 50 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to yield a crude product. The product was isolated by silica-gel flash chromatography using 10% ethylacetate / hexane as the eluting solvent to yield 2.1 g of pure product **11B**.

**Step 2:** Product **11B** from step 1 was dissolved in 5 mL toluene and NaOBu<sup>t</sup> (0.11g, 0.00114 mol), morpholine (0.092 mL, 0.001 mol), Pd (dba)<sub>2</sub> (0.024 g, 0.000026 mol), and P(*o*-tolyl)<sub>3</sub> (0.016g, 0.00005 mol) were added and the resulting




mixture was heated to 100°C for 24h. The reaction was cooled to room temperature and diluted with ethylacetate (50 mL) and washed with water and brine. The organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give a brown oil. The crude product was subjected to silica gel preparatory plate chromatography to yield 0.24 g of pure product **11C**.

**Step 3:** Product **11C** from **step 2** (0.24 g, 0.00069 mol) was dissolved in 10 mL anhydrous THF and cooled to -70°C and treated with n BuLi (2.5 M, 0.225 mL, 0.00055 mol). The mixture was stirred for 20 min and then a 5 mL anhydrous THF solution of pyridine disulfide was stirred and the resulting mixture was stirred for 2 h and allowed to warm to 0°C. The reaction was quenched with aq. ammonium chloride and diluted with 50 mL ethyl acetate. The organic phases were washed with water, brine (1 x 50 mL) dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to yield a crude product which was purified by flash chromatography with 5 to 20% ethylacetate/hexanes as the eluting solvent. Two regioisomeric products were obtained. The first fraction yielded 0.09g of product **13B**. The second fraction yielded regioisomeric product **13C** 0.1 g.

**Step 4:** The product **13B** (0.066 g, 0.000145 mol) from **step 3** was dissolved in 10 mL CH<sub>2</sub>Cl<sub>2</sub> and treated with mCPBA (70%, 0.095 g , 0.00036mol). The reaction was stirred for 15 h. The reaction was diluted with 50 mL CH<sub>2</sub>Cl<sub>2</sub> and washed with aq. NaHSO<sub>3</sub> (1x 25 mL), aq. NaHCO<sub>3</sub> (1 x 25 mL), and brine (1 x 25 mL). The organic phases were dried, filtered, and concentrated to yield a crude product that was purified by silica gel preparatory plate using ethyl acetate as the developing solvent to yield: 0.03g of **12G**.

Similarly, Product **13C** was oxidized with mCPBA to provide the corresponding sulfone **12H**.

# Synthesis of Compounds 6F and 7A



**Step 1:** NaOH (50 g) was dissolved in H<sub>2</sub>O (50 mL). Toluene (100 mL) was added, followed by addition of indole (I) (20 g, 0.17 mol). The resulting mixture was stirred at room temperature for 10 min. 2-Fluorophenylsulphonyl chloride (33 g, 0.17 mol), and tetrabutylammonium hydrogensulfate (cat.) were added. The reaction mixture was stirred at room temperature for 2 h. The aqueous layer was then removed, and the organic layer was washed with brine (3 × 100 mL). The resulting organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to dryness. The crude product was recrystallized using EtOAc (20 mL) and hexanes (70 mL) to give pure compound **3B** 40.5 g (86%) as white crystals.


**Step 2:** Product **3B** from step 1 (0.462 g, 0.0017 mol) was dissolved in THF (10 mL) and cooled to -78°C and treated with nBuLi (1.1 eq.). The resulting mixture was stirred for 1.5 h and then isopropylbenzaldehyde **4B** (5 mL THF solution) was added and stirred for additional 0.5h. The reaction was quenched with aq. ammonium chloride (10 mL) and allowed to warm to room temperature. The product was isolated with ethyl acetate (100 mL). The organics were dried and concentrated to yield a crude product that was purified by silica-gel chromatography using 9:1 hexane:ethylacetate as the eluting solvent to provide 0.486g of pure product **5B**.

**Step 3:** The product **5B** from Step 2 (0.1g, 0.0002mol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was treated in sequence with triethylsilane (0.38 mL, 10 eq.) and BF<sub>3</sub>·OEt. The mixture was stirred at room temperature for 1 h and then washed with aq. NaHCO<sub>3</sub> (10 mL) and brine (10 mL). The organics were dried and concentrated to yield a crude product

that was purified by silica gel preparatory plate chromatography to provide 0.025g of pure product **6F**.

**Step 4:** The product **5B** from **Step 2** (0.1g, 0.0002mol) in  $\text{CH}_2\text{Cl}_2$  (30 mL) was treated with celite (1g) and pyridiniumchlorochromate (PCC) (0.2g, 0.0008 mol) and the mixture was stirred overnight. The reaction was filtered and the filtrate was washed with aq.  $\text{NaHCO}_3$  (1 X 20 mL) and brine (1 x 20 mL). The organic phases were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to yield a crude product that was purified by preparatory plate chromatography using 3:7 ethylacetate:hexane to give 0.026g of pure product **7A**.

### Synthesis of Compound 12I




**Step 1:** Indole (**I**) (10 g, 0.085 mol) was dissolved in methylene chloride (200 mL). DMAP (cat.) and butoxycarbonyl anhydride (18.7 g, 0.085 mol) was added and the reaction was stirred at room temperature for 10 h. The reaction mixture was washed with aqueous NaCl solution (100 mL). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to dryness. The crude product was purified via silica gel chromatography (25% EtOAc/hexanes) to give 18.6 g (100%) of compound as a light yellow oil.

**Step 2:** The product from **step 1** (1 g, 0.0046 mol) in anhydrous THF (15 mL) at  $-78^\circ\text{C}$  was treated with n-BuLi (1.1. eq.) and stirred for 45 min. To this solution was added a THF (5 mL) solution of isopropylbenzenesulfonyl fluoride (1 g, 0.05mol). The resulting mixture was stirred for 4 h and the reaction mixture was allowed to warm to room temperature. The reaction was quenched with aq. NaCl. The organic phases were dried and concentrated to give a crude product, which was purified by silica gel

column chromatography using 5:1 hexane:ethylacetate as the eluting solvent. The fractions were collected and concentrated to yield pure product **11D** (0.466 g).

**Step 3:** The product **11D** from **step 2** (0.143 g, 0.00048 mol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was treated with NaOH (1 N, 5 mL) and catalytic amounts of tetrabutylammonium hydrogensulfate). To this mixture, pyridylsulfonyl chloride (0.1 g, 0.0006 mol) was added. The reaction was diluted with  $\text{CH}_2\text{Cl}_2$  (50 mL) and washed with water (1x 50 mL) and brine (1x 50 mL). The organic phases were dried and concentrated to yield a crude product that was purified by chromatography using 4:1 hexane: ethylacetate as the solvent to provide 0.089 g of pure product **12I**.

## Synthesis of Compounds 12J, 12K and 13D



**Step 1:** NaH (0.22g, 95% dry, 8.6 mmol) was dissolved in DMF (20 mL) followed by addition of indole (I) (1 g, 8.5 mmol). The mixture was stirred at room temperature for 10 min. Bis-2-pyridine disulfide (II) (2.17 g, 8.5 mmol) was added as a DMF solution. The reaction mixture was stirred at room temperature for 10 h. The reaction mixture was partitioned between EtOAc and water. The aqueous layer was extracted with additional EtOAc. The combined organic layer was washed with brine, then dried with  $\text{Na}_2\text{SO}_4$  and concentrated to dryness. The crude product was purified by silica gel chromatography (20% EtOAc / hexanes) to give 1.16 g of pure product as a white solid (III).

**Step 2:** Product from step 1 (0.76 g, 3.3 mmol) and  $\text{Boc}_2\text{O}$  (0.88 g, 4 mmol) were dissolved in methylene chloride (20 mL). DMAP (cat.) was added and the reaction was stirred at room temperature for 4 h. The reaction mixture was washed with aqueous NaCl solution (100 mL). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to give 1.1 g (100%) of the product as a light yellow solid 3C.


**Step 3:** In a flame-dried flask under  $\text{N}_2$  blanket, product 3C from step 2 (0.4 g, 1.2 mmol) was dissolved in dry THF (20 mL) and cooled to  $-78^\circ\text{C}$ . A solution of n-butyl lithium (1.9 M in hexanes, 0.75 mL, 1.4 mmol) was added and the reaction mixture was stirred for 45 min. Isopropylbenzenedisulfide (0.45 g, 1.0 mmol) in THF (1.5 mmol) was added and the reaction mixture was stirred at  $-78^\circ\text{C}$  for several hours before it was slowly warmed up to room temperature. The reaction mixture was then quenched with saturated aqueous  $\text{NH}_4\text{Cl}$  (20 mL). Ethyl acetate (30 mL) was added

and the layers were separated. The organic layer was washed with brine, then dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to dryness. The crude product was purified via silica gel chromatography (11% EtOAc/hexanes) to give 0.135 g (30%) of pure product **13D** as a colorless solid.

5       **Step 4:** Product **13D** from **step 3** (115 mg, 0.3 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (15 mL) and cooled to  $0^\circ\text{C}$  in an ice bath. mCPBA (410 mg, ca. 1.7 mmol) was added. The ice bath was removed and the reaction mixture was stirred at room temperature overnight. Aqueous  $\text{NaHCO}_3$  (200 mL) and  $\text{CH}_2\text{Cl}_2$  were added and the layers were separated. The organic layer was washed with aq  $\text{NaHSO}_3$ ,  $\text{NaHCO}_3$ ,  
10    $\text{H}_2\text{O}$ , and brine, then dried with  $\text{Na}_2\text{SO}_4$ . The crude product was purified by silica gel chromatography (33% EtOAc/hexanes) to give 78 mg (58%) of the product **12J** as a white solid.

**Step 5:** NaH (50 mg, 95% dry, 2 mmol) was dissolved in DMF (10 mL) at  $0^\circ\text{C}$  followed by addition of product **12J** from **step 4** (35 mg, 0.08 mmol). The mixture was  
15   stirred at  $0^\circ\text{C}$  for 10 min.  $\text{CH}_3\text{I}$  (excess, 1 mL) was added. The reaction mixture was slowly warmed up to room temperature and stirred for 2 h. The reaction mixture was partitioned between EtOAc and water. The aqueous layer was extracted with additional EtOAc (1 X 50 mL). The combined organic layers were washed with brine, then dried with  $\text{Na}_2\text{SO}_4$  and concentrated to dryness. The crude product was purified  
20   via silica gel chromatography (33% EtOAc / hexanes) to give 0.34 g (94%) of pure product **12K** as a white solid.

### Synthesis of Compound 12M




**Step 1:** In a flame-dried flask under N<sub>2</sub> blanket, 3-bromofuran (1.3 g, 9.0 mmol) was dissolved in dry THF (50 mL) and cooled to -78 °C. A solution of n-butyl lithium (1.6 M in hexanes, 5.6 mL, 9.0 mmol) was added and the reaction mixture was stirred for 15 min. Isopropylbenzenesulphonyl fluoride (i) (0.90 g, 3.0 mmol) in THF (3 mL) was added to the reaction mixture at -78 °C and stirred for 3 h. The reaction mixture was slowly warmed to room temperature and quenched with saturated aqueous NH<sub>4</sub>Cl (20 mL). Ethyl acetate (100 mL) was added and the layers were separated. The organic layer was washed with brine, then dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to dryness. The crude product was purified via prep. TLC (50% EtOAc/hexanes) to give 0.44 g (42%) of pure product **11E** as a white solid.

**Step 2:** In a flame-dried flask under N<sub>2</sub> blanket, compound **11E** (0.44 g, 1.3 mmol) was dissolved in dry THF (20 mL) and cooled to -78 °C. A solution of *t*-butyl lithium (1.5 M in hexanes, 1.7 mL, 2.5 mmol) was added and the reaction mixture was stirred for 15 min. 2-Fluorophenyl disulphide (0.32 g, 1.3 mmol) in THF (5 mL) was added and the reaction mixture was stirred at -78 °C for 1 h before being slowly warmed to 0 °C. The reaction mixture was then quenched with saturated aqueous NH<sub>4</sub>Cl (50 mL). Ethyl acetate (100 mL) was added and the layers were separated. The organic layer was washed with brine, then dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to dryness. The crude product **13E** was used without further purification.

**Step 3:** Compound **13E** (270 mg, 0.57 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (15 mL). mCPBA (210 mg, 57-86%, 0.69 mmol) was added and the solution was stirred at room temperature for 1 h. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL), washed with NaHSO<sub>3</sub> (40 mL) and NaHCO<sub>3</sub> (50 mL). The organic layers were combined and dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The crude material was purified via silica gel chromatography (50% EtOAc / hexanes) to give 120 mg (43%) of compound **12L** as a light yellow oil.

**Step 4:** Compound **12L** (70 mg, 0.14 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 mL). mCPBA (30 mg, 57-86%, 0.10 mmol) was added and the solution was stirred at room temperature, overnight. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL), washed with NaHSO<sub>3</sub> (40 mL) and NaHCO<sub>3</sub> (50 mL). The organic layers were combined and dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The crude material was purified via silica gel chromatography (50% EtOAc / hexanes) to give 0.59 g (82%) of **12M** as a white solid.

# Synthesis of Compound 12N



5

**Step 1:** A mixture of 2,4-dibromophenyl 1 (5 g, 0.02 mol), 1,2 dibromoethane (37 mL, 0.4 mol), aq. NaOH (14 mL, 3N) Bu<sub>4</sub>N<sup>+</sup>HSO<sub>4</sub><sup>-</sup> (0.34 g, 1 mmol) was stirred vigorously at 70 °C for 10 h. The mixture was cooled to room temperature. CH<sub>2</sub>Cl<sub>2</sub> (100 mL) was added. The organic layer was washed with NaOH (1N), HCl (1N), water and brine, respectively. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to dryness. The crude product was purified by recrystallization from ether:hexane to give 3.2 g (45%) of pure compound II as a white solid.


**Step 2:** In a flame-dried flask under N<sub>2</sub> blanket, compound II (1 g, 0.003 mol) was dissolved in dry THF (40 mL) and cooled to -100 °C. A solution of n-butyl lithium (2.5 M in hexanes, 1.32 mL) was added and the reaction mixture was stirred for 30 min. The reaction mixture was warmed to -78 °C. An additional equivalent of n-BuLi (2.5 M in hexanes, 1.32 mL) was added and the reaction mixture was stirred for 30 min followed by addition of MeLi (1.4 M, 5.6 mL, 7.8 mmol). To the resulting mixture was added isopropylbenzene sulfonyl fluoride (0.72 g, 0.0036 mol) in THF (5 mL) and the reaction mixture was stirred for 2 h before quenching with aq NH<sub>4</sub>Cl. The reaction was diluted with EtOAc and extracted with water (100 mL) and brine (50 mL), respectively. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to dryness. The crude product was purified via preparative thin layer chromatography (40% EtOAc/hexanes) to give 0.98g (38%) of compound 11F as a white solid.



**Step 3:** In a flame-dried flask under N<sub>2</sub> blanket, compound **11F** (0.3 g, 0.001 mol) was dissolved in dry THF (20 mL) and cooled to -78 °C. A solution of n-butyl lithium (1.7 M in hexanes, 0.476 mL) was added and the reaction mixture was stirred for 45 min. Pyridine disulphide (0.248 g, 1.25 mmol) in THF (5 mL) was added and the reaction mixture was stirred at -78 °C for 2 h before being slowly warmed to room temperature. The reaction mixture was then quenched with saturated aqueous NH<sub>4</sub>Cl (50 mL). Ethyl acetate (100 mL) was added and the aqueous and organic layers were separated. The organic layer was washed with brine, then dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to dryness. The crude product was purified via silica gel chromatography (25% EtOAc/hexanes) to give 0.28 g (48%) of compound **13F** as a white powder.

**Step 4:** Compound **13F** (0.28 g, 0.7 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) with trifluoroacetic acid (0.2 mL). mCPBA (0.358 g, 57-86%) was added and the solution was stirred at room temperature, overnight. The reaction mixture was taken up in CH<sub>2</sub>Cl<sub>2</sub>, washed with NaHSO<sub>3</sub> (40 mL) and NaHCO<sub>3</sub> (50 mL). The organic layers were combined and dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The crude material was purified via silica gel chromatography (50% EtOAC / hexanes) to give 0.09 g (42%) of compound **12N** as a white foam.

#### Synthesis of 38A



**Step1:** A mixture of 1,1,2 trichloroethane **31** (10mL), isopropyl thiophenol (2.19g, 0.014 mol), 1N NaOH (16 mL, 0.016 mol), and Aliquat® (tricaprylylmethyl

ammonium chloride; 0.01g; catalytic) were stirred for 4 days. The reaction was diluted with 50 mL of CH<sub>3</sub>OH and cooled to 0°C and then H<sub>2</sub>O<sub>2</sub> was added dropwise (14 mL, 30%) followed by ammoniummolybdate (0.1g, catalytic). The mixture was stirred vigorously for 24h. TLC analysis showed the formation of two products. The reaction  
5 was diluted with 100 mL ethylacetate and washed with 2 X 50 mL of water and 1 X 50 mL of brine. The organic phases were dried and concentrated to provide a crude product which was subjected to flash column chromatography using 10% ethylacetate: hexane as the eluting solvent. Appropriate fractions were collected and concentrated to provide 1.1g ( 4 mmol) of half-way oxidized sulfoxide product which was dissolved  
10 in 50 mL methylene chloride at 0°C and treated with metachloroperoxybenzoic acid (0.776g, 4.5 mmol) and stirred for 2 days. TLC analysis showed complete conversion to the desired sulfone **33A**. The solvent was removed under reduced pressure and the product redissolved in ethylacetate (100 mL). The organic phases were washed with aq. NaHCO<sub>3</sub> (4x 50 mL) and brine (1 x 50 mL), dried (MgSO<sub>4</sub>) and concentrated to  
15 provide pure product **33A**.

**Step2:** Product **33A** from step 1 (1.1 g, 4 mmol) in 50 mL CH<sub>2</sub>Cl<sub>2</sub> was cooled to 0°C and treated with triethylamine (0.6 mL, 4.4 mmol). The mixture was stirred for 1h. The solvent was removed and the crude product was redissolved in 100 mL ethylacetate and washed with water (2 X 50mL) and brine (1 X 50 mL), dried, and  
20 concentrated to yield pure product **34A**.










**Step 3:** 2-Fluorobenzene thiol (0.57 g, 4.4 mmol) in anhydrous THF (25 mL) at 0°C was treated with NaH (0.17 g, 60 % oil dispersion, 4.4. mmol). The mixture was stirred for 10 min and then chlorovinyl sulfone **34A** (0.9 g, 3.7 mmol) was added as a 5 mL THF solution. The mixture was allowed to warm up to room temperature and  
25 stirring was continued for 10h. The reaction was cooled to 0°C and quenched with water (5 mL). The resulting mixture was diluted with ethyl acetate (100 mL) and the organics were washed with water (2X 50 mL) and brine (1 X 50 mL). The organics were dried and concentrated to yield a crude product, which was subjected to silica gel chromatography using 10% ethylacetate/hexane as the eluting solvent. The  
30 fractions were collected and concentrated to yield pure product **36A**.

**Step 4:** Product **36A** from step 3 (0.28g, 0.8 mmol) was dissolved in glacial acetic acid (10 mL), cooled to 0°C and treated with H<sub>2</sub>O<sub>2</sub> (3 mL, 30% water) and conc. H<sub>2</sub>SO<sub>4</sub> (2 mL). The mixture was allowed to warm to room temperature and heated to

60°C for 3h. The reaction was cooled to room temperature and diluted with 50 mL ethylacetate. The organic phase was washed with water (3 X 50 mL) and brine (1 X 50 mL). The organic phases were dried and concentrated to yield a solid product, which was recrystallized from ether as white solid **37A**.

- 5 **Step 5:** Product **37A** from step 4 (0.15g, 0.4 mmol) and furan (IV) (0.15 g, 2 mmol) in 2 mL chloroform were stirred for 10 h. The solvent was removed and the addition of CH<sub>3</sub>OH (5 mL) led to precipitation of product **38A** as a white solid. The product was a mixture of inseparable stereoisomers.

10 Table 2: Spectral Data for selected Compounds in Table 1

Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
A	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> -			-CF <sub>3</sub>	F	435.1(M-1, 100), 436.1(M, 50), 459(M+23, 12)
B	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> -			-CF <sub>3</sub>		603(M+1, 100), 547(63)
C	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> -			-OCF <sub>3</sub>	F	453(M+1, 13.54), 470.1(74), 475(85)
D	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> -			-OCF <sub>3</sub>	-NH(CH <sub>2</sub> ) <sub>2</sub> OH	453(M+1, 13.54), 470.1(74), 475(85)

Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
E	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	-CH <sub>3</sub>	489(M+1, 47), 506(16), 511[M+23, (30)], 548(13)
F	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	F	503(M+1, 67), 504(22, 64), 525(M+23, 100)
G	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	H	486 (M+1, 100), 508[M+23, (35.08)]
H	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	F	458 (M+1, 100), 183 (90)
I	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCF <sub>3</sub>	H	487.1(M+1, 100), 509[M+23, (80)], 425(75)
J	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> -			H	F	406 (M+1, 80), 275 (100)
K		-S(O <sub>2</sub> )-			Cl	F	451.1(M+1, 100), 130.0 (60)
L	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H	441 (M+1, 100), 377(10)







Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
M	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-N(CH <sub>3</sub> ) <sub>2</sub>	F	462(M+1, 100), 484[M+ 23, (30)]
N	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H	444 (M+1, 100)
O	-N(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	445(M+1, 100)
P	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-N(CH <sub>3</sub> ) <sub>2</sub>	H	445.1(M+1, 100), 431(10)
Q	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	460.1(M+1, 100), 482(M+23, (50))
R	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OH	F	435(M+1, 100), 457(M+ 23 (10))
S	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-				F	493(M+1, 100), 515(M+23(10))

Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
T	-CH(CH <sub>3</sub> ) <sub>2</sub>	covalent bond			-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>3</sub>	464[(M-HCl)+1, 100]
U	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-				H	487(M+1, 100)
V		-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	487(M+1, 100)
W		-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	H	464(M+1, 100)
X	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	472(M+1, 100), 486(50), 504(45)
Y	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	474.1(M+1, 100)
Z		-S(O <sub>2</sub> )-				H	466.1(M+1, 100), 488(40)
AA		-S(O <sub>2</sub> )-				H	455(100), 477(50), 346(30), 232(75), 200(80)
AB	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CN	H	427.1(M+1, 100), 253 (12), 149.0 (17)

Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
AC	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-				H	460 (M+1, 100)
AD	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CF <sub>3</sub>	H	470 (M+1, 50), 492 (50), 134 (30)
AE	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	F	437 (M+1, 369, 277)
AF	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	F	449 (M+1, 65), 471 (100), 159 (65)
AG	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	H	454.1 (M+1, 100), 432 (50)
AH		-S(O <sub>2</sub> )-			-OCH <sub>3</sub>	F	447.1 (M+1, 100), 329.1 (10)
AI		-S(O <sub>2</sub> )-				H	440.1 (M+1, 100), 246.0 (6), 185.0 (6)

Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
AJ	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	444(M+1)
AK	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH(CH <sub>3</sub> ) <sub>2</sub>	F	477(M+1), 357(10)
AL	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH(CH <sub>3</sub> ) <sub>2</sub>	H	460.1(M+1, 100), 418(05)
AM	-OCH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CH(CH <sub>3</sub> ) <sub>2</sub>	H	460(M+1, 100), 482(M+2 3, (25))
AN	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-OCH(CH <sub>3</sub> ) <sub>2</sub>	-C(O)OCH <sub>3</sub>	518(M+1, 100), 540(50), 248(75), 210(65)
AO	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	F	(472 (M+1, 100), 376 (30)
AP	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H	441, (M+1, 100)




Cmpd.	R <sup>1</sup>	L <sup>1</sup>	M <sup>1</sup> -Y	M <sup>2</sup> -Z	Y	Z	MS
AQ	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H	(455 (M+1, 100)
AR	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			-CF <sub>3</sub>	H	475.1(M+ 1, 100), 497(M+2 3, (45)), 459(35)
AS	-CH(CH <sub>3</sub> ) <sub>2</sub>	-S(O <sub>2</sub> )-			H	H	392 (M+1)

It will be understood that various modifications may be made to the  
embodiments and examples disclosed herein. Therefore, the above description  
should not be construed as limiting, but merely as exemplifications of preferred  
embodiments. Those skilled in the art will envision various modifications within the  
scope and spirit of the claims appended hereto.

We claim:

1. A compound of formula I



or a pharmaceutically acceptable salt or solvate thereof, wherein

5 n is 0 to 4;

p is 0 to 4;

q is 0 to 5;

X is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, haloalkoxy, haloalkyl, halogen, heteroalkyl, heteroaryl, -CF<sub>3</sub>, -CN, -C(O)N(R<sup>2</sup>)<sub>2</sub>, -C(O)OR<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub>, -NHC(O)R<sup>2</sup>,  
 10 -NR<sup>2</sup>C(O)OR<sup>2</sup>, -NR<sup>2</sup>C(O)N(R<sup>2</sup>)<sub>2</sub>, -NO<sub>2</sub>, -NC(=N-CN)NHR<sup>2</sup>, -OCF<sub>2</sub>H, -OCF<sub>3</sub>, -OH and -S(O<sub>2</sub>)N(R<sup>2</sup>)<sub>2</sub>, with the proviso that when n is 2, 3 or 4, the X moieties can be the same or different and are independently selected from the group listed above;

Y is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, halogen, heteroalkyl, heteroaryl,  
 15 -O-cycloalkyl, -CF<sub>3</sub>, -CN, -C(O)OR<sup>2</sup>, -C(O)R<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -OCF<sub>2</sub>H and -OH, with the proviso that:

when p is 2, 3 or 4, the Y moieties can be the same or different and are independently selected from the group listed above; or

20 when p is 2, the Y moieties can form a cyclic ring of 3 to 7 ring atoms of which 1-2 may be a heteroatom;

Z is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl, cycloalkenyl, halogen, heteroalkyl, heteroaryl, heterocyclyl, heterocyclenyl, -O-cycloalkyl, -CF<sub>3</sub>, -CN, -C(O)OR<sup>2</sup>, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -OCF<sub>2</sub>H, -OH and -S(O<sub>2</sub>)R<sup>2</sup>, with  
 25 the proviso that when q is 2, 3, 4 or 5, the Z moieties can be the same or different and are independently selected from the group listed above;

R<sup>1</sup> is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl, heterocyclenyl, and -N(R<sup>2</sup>)<sub>2</sub>;

$R^2$  is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl, heterocyclenyl, and hydroxyalkyl;

$L^1$  is selected from the group consisting of a covalent bond,  $-C(F_2)-$ ,  $-(CH(OR^2))-$ ,  $-C(O)-$ ,  $-C(O)N(H)-$ ,  $-C(=N-OR^2)-$ ,  $-C(=NR^2)-$ ,  $-C(=N-CN)-$ ,  $-C(R^2)_2-$ ,  $-N(R^2)-$ ,  $-N(H)C(O)-$ ,  $-N(R^2)S(O_2)-$ ,  $-O-$ ,  $-OC(O)-$ ,  $-C(O)O-$ ,  $-S-$ ,  $-S(O_2)-$ ,  $-S(O)-$  and  $-S(O_2)N(R^2)-$ , with the proviso that when  $L^1$  is a covalent bond,  $M^1$  is directly attached to the phenyl carbon marked A;

$L^2$  is selected from the group consisting of a covalent bond,  $-C(R^2)_2-$ ,  $-C(=N-OR^2)-$ ,  $-C(O)-$ ,  $-C(O)N(H)-$ ,  $-C(O)O-$ ,  $-OC(O)-$ ,  $-N(H)C(O)-$ ,  $-NHS(O_2)-$ ,  $-N(R^2)-$ ,  $-O-$ ,  $-S-$ ,  $-S(O)-$ ,  $-S(O_2)-$  and  $-S(O_2)N(R^2)-$ , with the proviso that when  $L^2$  is a covalent bond,  $M^1$  is directly attached to  $M^2$ ;

$M^1$  is aryl, cycloalkyl, heteroaryl or heterocycloalkyl;  
and

$M^2$  is alkyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclyl, heterocyclenyl,  $-C(O)R^2$ ,  $-C(O)OR^2$ ,  $-N(R^2)_2$  or  $-S(O_2)R^2$ ;  
with the provisos that:

- i) when  $M^2$  is  $-N(R^2)_2$ ,  $L^2$  is a covalent bond,  $-CH_2-$ ,  $-C(O)-$  or  $-S(O_2)-$  and Z is absent;
  - ii) when  $M^2$  is  $-C(O)R^2$  or  $-C(O)OR^2$ ,  $L^2$  is a covalent bond,  $-CH_2-$ ,  $-NH-$  or  $-N(alkyl)-$  and Z is absent;
  - iii) when  $M^2$  is  $-S(O_2)R^2$ ,  $L^2$  is a covalent bond,  $-CH_2-$ ,  $-NH-$  or  $-N(alkyl)-$  and Z is absent;
- and
- iv) the two  $R^2$  moieties of  $-N(R^2)_2$  and  $-C(R^2)_2-$  are the same or different and are independently selected, or the two  $R^2$  of  $-N(R^2)_2$  are joined together and with the nitrogen to which they are attached to form a heterocyclic ring having 3 to 7 ring atoms optionally containing additionally one or more N or O atoms wherein said additional N can be optionally substituted with  $R^2$ ;

wherein each of said alkoxy, alkyl, aralkyl, aryl, cycloalkyl, cycloalkenyl, heteroaralkyl, heteroaryl, heterocyclyl, and heterocyclenyl in the definitions above can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen,  $-CF_3$ , alkoxy,  $-CN$ ,  $-C(O)N(R^2)_2$ ,  $-C(O)OR^2$ ,  $-C(O)R^2$ ,

-NC(O)R<sup>2</sup>, -NR<sup>2</sup>C(O)OR<sup>2</sup>, -NR<sup>2</sup>C(O)N(R<sup>2</sup>)<sub>2</sub>, -NC(=N-CN)NHR<sup>2</sup>, -NO<sub>2</sub>, -N(R<sup>2</sup>)<sub>2</sub>,  
-OCF<sub>2</sub>H, -OCF<sub>3</sub>, -OH, -S(O<sub>2</sub>)R<sup>2</sup> and -S(O<sub>2</sub>)N(R<sup>2</sup>)<sub>2</sub>.

2. The compound according to claim 1, or a pharmaceutically acceptable salt or  
5 solvate thereof, wherein

M<sup>1</sup> is aryl, heteroaryl or heterocycloalkyl;

and

M<sup>2</sup> is aryl, cycloalkyl, heteroaryl, or heterocyclyl.

- 10 3. The compound according to claim 1, or a pharmaceutically acceptable salt or  
solvate thereof, wherein X is selected from the group consisting of hydrogen, alkoxy,  
-CF<sub>3</sub>, haloalkoxy, halogen, -OCF<sub>3</sub>, -OCF<sub>2</sub>H and -OH.

4. The compound according to claim 1, or a pharmaceutically acceptable salt or  
15 solvate thereof, wherein Y is selected from the group consisting of hydrogen, alkoxy,  
alkyl, -CF<sub>3</sub>, -C(O)OR<sup>2</sup>, cycloalkyl, halogen, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -O-cycloalkyl and -OH.

5. The compound according to claim 1, or a pharmaceutically acceptable salt or  
solvate thereof, wherein Z is selected from the group consisting of hydrogen, alkoxy,  
20 alkyl, -CF<sub>3</sub>, -C(O)OR<sup>2</sup>, halogen, heterocyclyl, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub>, -O-cycloalkyl and -OH.

6. The compound according to claim 1, or a pharmaceutically acceptable salt or  
solvate thereof, wherein R<sup>1</sup> is selected from the group consisting of hydrogen, alkoxy,  
alkyl, cycloalkyl and -N(R<sup>2</sup>)<sub>2</sub>.

25

7. The compound according to claim 1, or a pharmaceutically acceptable salt or  
solvate thereof, wherein R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl,  
aryl, cycloalkyl and heterocycloalkyl.

- 30 8. The compound according to claim 1, or a pharmaceutically acceptable salt or  
solvate thereof, wherein L<sup>1</sup> is selected from the group consisting of -C(R<sup>2</sup>)<sub>2</sub>-,  
-N(R<sup>2</sup>)S(O<sub>2</sub>)-, -N(R<sup>2</sup>)-, -S(O<sub>2</sub>)- and -S(O<sub>2</sub>)N(R<sup>2</sup>)-.

9. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, wherein  $L^2$  is selected from the group consisting of a covalent bond,  $-C(R^2)_2-$ ,  $-N(R^2)S(O_2)-$ ,  $-N(R^2)-$ ,  $-S(O_2)-$  and  $-S(O_2)N(R^2)-$ .
10. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, wherein  $M^1$  is aryl, heteroaryl or heterocyclyl.
11. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, wherein  $M^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl.
12. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, wherein
- X is selected from the group consisting of hydrogen, alkoxy,  $-CF_3$ , haloalkoxy, halogen,  $-OCF_3$ ,  $-OCF_2H$  and  $-OH$ ;
- Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ ,  $-C(O)OR^2$ , cycloalkyl, halogen,  $-N(R^2)_2$ ,  $-OCF_3$ ,  $-O$ -cycloalkyl and  $-OH$ ;
- Z is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-CF_3$ ,  $-C(O)OR^2$ , halogen, heterocyclyl,  $-N(R^2)_2$ ,  $-OCF_3$ ,  $-O$ -cycloalkyl and  $-OH$ ;
- $R^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-N(R^2)_2$ ;
- $R^2$  is selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl and heterocyclyl;
- $L^1$  is selected from the group consisting of  $-C(R^2)_2-$ ,  $-N(R^2)-$ ,  $-S(O_2)-$  and  $-S(O_2)N(R^2)-$ ;
- $L^2$  is selected from the group consisting of a covalent bond,  $-C(R^2)_2-$ ,  $-N(R^2)S(O_2)-$ ,  $-N(R^2)-$ ,  $-S(O_2)-$  and  $-S(O_2)N(R^2)-$ ;
- $M^1$  is aryl, heteroaryl or heterocyclyl;
- $M^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl;
- with the provisos that:
- the two  $R^2$  moieties of  $N(R^2)_2$  and  $C(R^2)_2-$  are the same or different and are independently selected, wherein each  $R^2$  of  $N(R^2)_2$  are joined together and with the nitrogen to which they are attached form a heterocyclic ring having 3 to 7 ring atoms;

wherein each of said alkoxy, alkyl, aralkyl, aryl, cycloalkyl, heteroaralkyl, heteroaryl and heterocyclyl in the definitions above can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of

5 halogen, alkoxy,  $-\text{CF}_3$ ,  $-\text{CN}$ ,  $-\text{C}(\text{O})\text{N}(\text{R}^2)_2$ ,  $-\text{C}(\text{O})\text{OR}^2$ ,  $-\text{C}(\text{O})\text{R}^2$ ,  $-\text{NC}(\text{O})\text{R}^2$ ,  $-\text{NR}^2\text{C}(\text{O})\text{OR}^2$ ,  $-\text{NR}^2\text{C}(\text{O})\text{N}(\text{R}^2)_2$ ,  $-\text{NC}(=\text{N}-\text{CN})\text{NHR}^2$ ,  $-\text{NO}_2$ ,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_2\text{H}$ ,  $-\text{OCF}_3$ ,  $-\text{OH}$ ,  $-\text{S}(\text{O}_2)\text{R}^2$  and  $-\text{S}(\text{O}_2)\text{N}(\text{R}^2)_2$ .

13. The compound according to claim 1 or a pharmaceutically acceptable salt or  
10 solvate thereof, wherein

n is 1;

X is hydrogen and  $-\text{OH}$ ;

Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , cycloalkyl, halogen,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$  and  $-\text{OH}$ ;

15 Z is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , halogen, heterocyclyl,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$ ,  $-\text{O-cycloalkyl}$  and  $-\text{OH}$ ;

$\text{R}^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-\text{N}(\text{R}^2)_2$ ;

$\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl and  
20 heterocyclyl;

$\text{L}^1$  is selected from the group consisting of  $-\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^2)_2-$  and  $-\text{S}(\text{O}_2)-$ ;

$\text{L}^2$  is selected from the group consisting of a covalent bond,  $-\text{C}(\text{R}^2)_2-$ ,  $-\text{C}(\text{O})-$  and  $-\text{S}(\text{O}_2)-$ ;

$\text{M}^1$  is aryl, heteroaryl or heterocyclyl;

25 and

$\text{M}^2$  is aryl, cycloalkyl, heteroaryl, or heterocyclyl.

14. The compound according to claim 1 or a pharmaceutically acceptable salt or  
30 solvate thereof, wherein

n is 1;

X is hydrogen and  $-\text{OH}$ ;

Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{C}(\text{O})\text{OR}^2$ , cycloalkyl,  $-\text{CF}_3$ , halogen,  $-\text{OCF}_3$  and  $-\text{OH}$ ;

Z is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ , halogen,  $-\text{OCF}_3$ ,  $-\text{O-cycloalkyl}$  and  $-\text{OH}$ ;

$\text{R}^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and  $-\text{N}(\text{R}^2)_2$ ;

5  $\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, aryl and cycloalkyl;

$\text{L}^1$  and  $\text{L}^2$  are the same or different and are independently  $-\text{C}(\text{R}^2)_2-$  or  $-\text{S}(\text{O}_2)-$ ;


$\text{M}^1$  is aryl or heteroaryl;

and

$\text{M}^2$  is aryl, cycloalkyl, heteroaryl or heterocyclyl.

10

15. The compound according to claim 1 having the formula II:



or a pharmaceutically acceptable salt thereof, wherein

Y is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{CN}$ ,  $-\text{C}(\text{O})\text{OR}^2$ , cycloalkyl, halogen,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$  and  $-\text{OH}$ , with the proviso that when p is 2, the Y moieties can form a cyclic ring of 3 to 7 ring atoms of which 1-2 may be a heteroatom;

15

Z is selected from the group consisting of hydrogen, alkoxy, alkyl,  $-\text{CF}_3$ ,  $-\text{C}(\text{O})\text{OR}^2$ , halogen, heterocyclyl,  $-\text{N}(\text{R}^2)_2$ ,  $-\text{OCF}_3$ ,  $-\text{O-cycloalkyl}$  and  $-\text{OH}$ ;

20

$\text{R}^1$  is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl, heterocyclyl and  $-\text{N}(\text{R}^2)_2$ ;

$\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, aryl and cycloalkyl;

$\text{L}^1$  is a covalent bond,  $-\text{C}(\text{R}^2)_2-$  or  $-\text{S}(\text{O}_2)-$ ;












$\text{M}^1$  is aryl, indolyl, oxabicycloheptenyl or furanyl;

25














and















$\text{M}^2$  is aryl, cycloalkyl, heteroaryl or heterocyclyl.

16. The compound according to claim 15 having the formula II, or a pharmaceutically acceptable salt thereof, wherein  $R^1$ ,  $L^1$ ,  $M^1$ ,  $M^2$ , Y and Z are as set forth in the following table:




















	$R^1$	$L^1$	$M^1$ -Y	$M^2$ -Z	Y	Z
	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{CF}_3$	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{CF}_3$	
	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{OCF}_3$	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			$-\text{OCF}_3$	$-\text{NH}(\text{CH}_2)_2\text{OH}$
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCF}_3$	$-\text{CH}_3$
































	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	$Y$	$Z$
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCF}_3$	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCF}_3$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCF}_3$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2-$			H	F
		$-\text{S}(\text{O}_2)-$			Cl	F

	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	Y	Z
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$\text{N}(\text{CH}_3)_2$	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	H
	$-\text{N}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{N}(\text{CH}_3)_2$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			OH	F

	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	Y	Z
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$				F
	$-\text{CH}(\text{CH}_3)_2$	Covalent bond			$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_3$
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$				H
		$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H
		$-\text{S}(\text{O}_2)-$			$-\text{OCH}_3$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H


	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	$Y$	$Z$
		$-S(O_2)-$				H
		$-S(O_2)-$				H
	$-CH(CH_3)_2$	$-S(O_2)-$			$-CN$	H
	$-CH(CH_3)_2$	$-S(O_2)-$				H
	$-CH(CH_3)_2$	$-S(O_2)-$			$-CF_3$	H
	$-CH(CH_3)_2$	$-S(O_2)-$			H	F
	$-CH(CH_3)_2$	$-S(O_2)-$			$-OCH_3$	F

	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	$Y$	$Z$
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCH}_3$	H
		$-\text{S}(\text{O}_2)-$			$-\text{OCH}_3$	F
		$-\text{S}(\text{O}_2)-$				H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCH}(\text{CH}_3)_2$	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCH}(\text{CH}_3)_2$	H
	$-\text{OCH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CH}(\text{CH}_3)_2$	H

	$R^1$	$L^1$	$M^1-Y$	$M^2-Z$	$Y$	$Z$
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{OCH}(\text{CH}_3)_2$	$-\text{COOCH}_3$
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	F
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			$-\text{CF}_3$	H
	$-\text{CH}(\text{CH}_3)_2$	$-\text{S}(\text{O}_2)-$			H	H

17. The compound according to claim 1 having the formula III:

78



or a pharmaceutically acceptable salt thereof, wherein


Y is selected from the group consisting of hydrogen, alkoxy, alkyl, -CF<sub>3</sub>, cycloalkyl, halogen, -OCF<sub>3</sub> and -OH;

5 Z is selected from the group consisting of hydrogen, alkyl, -CF<sub>3</sub>, halogen, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub> and -OH;

and

M<sup>2</sup> is aryl or heteroaryl.

10 18. The compound according to claim 1 having the formula IV



or a pharmaceutically acceptable salt thereof, wherein


Y is selected from the group consisting of hydrogen, alkoxy, alkyl, cycloalkyl and -OCF<sub>3</sub>;

15 Z is selected from the group consisting of hydrogen, alkyl, -CF<sub>3</sub>, halogen, -N(R<sup>2</sup>)<sub>2</sub>, -OCF<sub>3</sub> and -OH;

and

$M^2$  is aryl or heteroaryl.

19. The compound according to claim 1 having the formula V



5 or a pharmaceutically acceptable salt thereof, wherein


Z is selected from the group consisting of hydrogen,  $-CF_3$ , halogen,  $-OCF_3$  and  $-OH$ ;

and

$M^2$  is aryl or heteroaryl.

10

20. The compound of claim 1 having the formula VI



or a pharmaceutically acceptable salt thereof, wherein


$M^2$  is aryl or heteroaryl;

15 and

Z is selected from the group consisting of hydrogen, alkyl, halogen,  $-CF_3$ ,  $-N(R^2)_2$ ,  $-OH$  and  $-OCF_3$ .



21. The compound of claim 1 having the formula:



5 or a pharmaceutically acceptable salt thereof.

22. A pharmaceutical composition comprising an effective amount of at least one compound according to claim 1 and a pharmaceutically acceptable carrier.

10 23. A pharmaceutical composition comprising an effective amount of at least one compound according to claim 16 and a pharmaceutically acceptable carrier.

24. A method of treating cancer, inflammatory diseases, immunomodulatory diseases, or respiratory diseases comprising administering to a mammal in need of  
15 such treatment an effective amount of at least one compound according to claim 1.

25. A method of treating cancer, inflammatory diseases, immunomodulatory diseases, or respiratory diseases comprising administering to a mammal in need of such treatment an effective amount of at least one compound according to claim 16.
- 5 26. A method of treating cutaneous T cell lymphoma, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, sepsis, shock, sarcoidosis, idiopathic pulmonary fibrosis, bronchopulmonary dysplasia, retinal disease, scleroderma, osteoporosis, renal ischemia, myocardial infarction, cerebral stroke, cerebral ischemia, nephritis, hepatitis, glomerulonephritis, cryptogenic fibrosing  
10 aveolitis, psoriasis, transplant rejection, atopic dermatitis, vasculitis, allergy, seasonal allergic rhinitis, Crohn's disease, inflammatory bowel disease, reversible airway obstruction, adult respiratory distress syndrome, asthma, chronic obstructive pulmonary disease (COPD) or bronchitis comprising administering to a mammal in need of such treatment an effective amount of at least one compound according to  
15 claim 1.
27. A method of treating cutaneous T cell lymphoma, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, sepsis, shock, sarcoidosis, idiopathic pulmonary fibrosis, bronchopulmonary dysplasia, retinal  
20 disease, scleroderma, osteoporosis, renal ischemia, myocardial infarction, cerebral stroke, cerebral ischemia, nephritis, hepatitis, glomerulonephritis, cryptogenic fibrosing aveolitis, psoriasis, transplant rejection, atopic dermatitis, vasculitis, allergy, seasonal allergic rhinitis, Crohn's disease, inflammatory bowel disease, reversible airway obstruction, adult respiratory distress syndrome, asthma, chronic obstructive  
25 pulmonary disease (COPD) or bronchitis comprising administering to a mammal in need of such treatment an effective amount of at least one compound according to claim 16.
28. The method of claim 24 wherein the condition or disease treated is selected  
30 from rheumatoid arthritis, multiple sclerosis, seasonal allergic rhinitis, psoriasis, transplant rejection and chronic obstructive pulmonary disease.

29. The method of claim 25 wherein the condition or disease treated is selected from rheumatoid arthritis, multiple sclerosis, seasonal allergic rhinitis, psoriasis, transplant rejection and chronic obstructive pulmonary disease.

5 30. A process for making a pharmaceutical composition comprising combining at least one compound of claim 1 and at least one pharmaceutically acceptable carrier.

31. A process for making a pharmaceutical composition comprising combining at least one compound of claim 16 and at least one pharmaceutically acceptable carrier.

10

32. A method of treating rheumatoid arthritis comprising administering to a mammal in need thereof an effective amount of at least one compound of claim 1 in combination with at least one compound selected from the class consisting of a COX-2 inhibitor, a COX-1 inhibitor, an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of rheumatoid arthritis.

15

33. A method of treating rheumatoid arthritis comprising administering to a mammal in need thereof an effective amount of at least one compound of claim 16 in combination with at least one compound selected from the class consisting of a COX-2 inhibitor, a COX-1 inhibitor, an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of rheumatoid arthritis.

20

25 34. The method of claim 32 wherein the COX-2 inhibitor is Celebrex or Vioxx, the COX-1 inhibitor is Feldene, the immunosuppressive is methotrexate, leflunomide, sulfasalazine, or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.

35. The method of claim 33 wherein the COX-2 inhibitor is Celebrex or Vioxx, the COX-1 inhibitor is Feldene, the immunosuppressive is methotrexate, leflunomide, sulfasalazine, or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.

30

36. A composition for treating rheumatoid arthritis which comprises a compound selected from the class consisting of a COX-2 inhibitor, a COX-1 inhibitor, an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of rheumatoid arthritis and an effective amount of at least one compound of claim 1.
37. A composition for treating rheumatoid arthritis which comprises a compound selected from the class consisting of a COX-2 inhibitor, a COX-1 inhibitor, an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of rheumatoid arthritis and an effective amount of at least one compound of claim 16.
38. The composition of claim 36 wherein the COX-2 inhibitor is Celebrex or Vioxx, the COX-1 inhibitor is Feldene, the immunosuppressive is methotrexate, leflunomide, sulfasalazine, or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.
39. The composition of claim 37 wherein the COX-2 inhibitor is Celebrex or Vioxx, the COX-1 inhibitor is Feldene, the immunosuppressive is methotrexate, leflunomide, sulfasalazine, or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.
40. A method of treating multiple sclerosis comprising administering to a mammal in need thereof an effective amount of at least one compound of claim 1 in combination with an effective amount of a compound selected from Avonex, Betaseron, Copaxone or other compounds indicated for the treatment of multiple sclerosis.
41. A method of treating multiple sclerosis comprising administering to a mammal in need thereof an effective amount of at least one compound of claim 16 in combination with an effective amount of a compound selected from Avonex, Betaseron, Copaxone or other compounds indicated for the treatment of multiple sclerosis.

42. A composition for treating multiple sclerosis which comprises a compound selected from Avonex, Betaseron, Copaxone or other compounds indicated for the treatment of multiple sclerosis and an effective amount of at least one compound of claim 1.

5 43. A composition for treating multiple sclerosis which comprises a compound selected from Avonex, Betaseron, Copaxone or other compounds indicated for the treatment of multiple sclerosis and an effective amount of at least one compound of claim 16.

10 44. A method of treating psoriasis comprising administering to a mammal in need thereof an effective amount of at least one compound as defined in claim 1 in combination with a compound selected from the class consisting of an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of psoriasis.

15 45. A method of treating psoriasis comprising administering to a mammal in need thereof an effective amount of at least one compound as defined in claim 16 in combination with a compound selected from the class consisting of an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of psoriasis.

20 46. The method of claim 44 wherein the immunosuppressive is methotrexate, leflunomide, sulfasalazine or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.

25 47. The method of claim 45 wherein the immunosuppressive is methotrexate, leflunomide, sulfasalazine or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.

30 48. A composition for treating psoriasis which comprises a compound selected from the class consisting of an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of psoriasis and an effective amount of at least one compound of claim 1.

49. A composition for treating psoriasis which comprises a compound selected from the class consisting of an immunosuppressive, a steroid, an anti-TNF- $\alpha$  compound or other classes of compounds indicated for the treatment of psoriasis and an effective amount of at least one compound of claim 16.
50. The composition of claim 48 wherein the immunosuppressive is methotrexate, leflunomide, sulfasalazine or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.
51. The composition of claim 49 wherein the immunosuppressive is methotrexate, leflunomide, sulfasalazine or cyclosporin, the steroid is  $\beta$ -methasone and the anti-TNF- $\alpha$  compound is Enbrel or Remicade.
52. A method of treating seasonal allergic rhinitis and/or asthma comprising an effective amount of at least one compound of claim 1 in combination with at least one H1 antagonist.
53. A method of treating seasonal allergic rhinitis and/or asthma comprising an effective amount of at least one compound of claim 16 in combination with at least one H1 antagonist.
54. A composition for treating seasonal allergic rhinitis and/or asthma which comprises an effective amount of at least one H1 antagonist and an effective amount of at least one compound of claim 1.
55. A composition for treating seasonal allergic rhinitis and/or asthma which comprises an effective amount of at least one H1 antagonist and an effective amount of at least one compound of claim 16.
56. The composition of claim 54 wherein the H1 antagonist is selected from Claritin, Clarinex, Zyrtec and Allegra.

57. The composition of claim 55 wherein the H1 antagonist is selected from Claritin, Clarinex, Zyrtec and Allegra.

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
7 October 2004 (07.10.2004)

PCT

(10) International Publication Number  
**WO 2004/085385 A3**

(51) International Patent Classification<sup>7</sup>: C07C 317/14,  
317/22, 317/36, C07D 209/30, 211/54, 213/71, 239/38,  
295/205, 307/64, 307/82, 401/12, 405/12, 413/12, A61K  
31/10, 31/136

(21) International Application Number:  
PCT/US2004/008333

(22) International Filing Date: 18 March 2004 (18.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/456,268 20 March 2003 (20.03.2003) US

(71) Applicant (for all designated States except US): SCHER-  
ING CORPORATION [US/US]; 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SHANKAR,  
Bandarpalle, B. [US/US]; 1124 Van Arsdale Drive,  
Branchburg, NJ 08853 (US). RIZVI, Razia, K. [US/US];  
22 Tomar Court, Bloomfield, NJ 07003 (US). KO-  
ZLOWSKI, Joseph, A. [US/US]; 1066 Stuart Road,  
Princeton, NJ 08540 (US). SHIH, Neng-Yang [US/US]; 4  
Overlook Drive, Warren, NJ 07059 (US).

(74) Agent: BLINKA, Thomas, A.; Schering-Plough Corpora-  
tion, Patent Department, K-6-1 1990, 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,

TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-  
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,  
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,  
EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,  
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,  
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,  
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ,  
TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM,  
ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD,  
SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY,  
KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG,  
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT,  
LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF,  
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
TD, TG)

— as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations


#### Published:

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
25 November 2004

[Continued on next page]

(54) Title: CANNABINOID RECEPTOR LIGANDS



(57) Abstract: There are disclosed compounds of the  
formula (I) a pharmaceutically acceptable salt or solvate  
of the compound, which exhibit anti-inflammatory  
and immunomodulatory activity. Also disclosed are  
pharmaceutical compositions containing said compounds.





---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US2004/008333

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C317/14 C07C317/22 C07C317/36 C07D209/30 C07D211/54  
 C07D213/71 C07D239/38 C07D295/205 C07D307/64 C07D307/82  
 C07D401/12 C07D405/12 C07D413/12 A61K31/10 A61K31/136

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>US 6 100 259 A (ELLIOTT JOHN DUNCAN ET AL) 8 August 2000 (2000-08-08)</p> <p>column 2, lines 34-45; claim 1; examples 1-63</p> <p>----- -/--</p>	<p>1-14, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56</p>

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

27 August 2004

Date of mailing of the international search report

07/10/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Cooper, S

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/008333

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/060447 A (DENG HONGFENG ; MAKRIYANNIS ALEXANDROS (US); UNIV CONNECTICUT (US)) 8 August 2002 (2002-08-08)  claim 1; tables 1,2	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
X	US 5 925 768 A (BARTH FRANCIS ET AL) 20 July 1999 (1999-07-20)  claim 21; table 1	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
X	WO 01/28557 A (DENG HONGFENG ; MAKRIYANNIS ALEXANDROS (US); UNIV CONNECTICUT (US)) 26 April 2001 (2001-04-26)  page 18, lines 2-16; claims 1,6; tables 1,2	1-14,16, 18,20, 22,24, 26,28, 30,32, 34,36, 38,40, 42,44, 46,48, 50,52, 54,56
X	WO 01/32169 A (MECHOULAM RAPHAEL ; TCHILIBON SUSANNA (IL); YISSUM RES DEV CO (IL); FR) 10 May 2001 (2001-05-10)  claims 1,9,17	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56

-/--

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/008333

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 03/020217 A (LIU QIAN ; MAKRIYANNIS ALEXANDROS (US); UNIV CONNECTICUT (US)) 13 March 2003 (2003-03-13)</p> <p>Intermediates A-E,G,H, claims 1,32-35; examples 1-6</p>	<p>1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56</p>
X	<p>EP 0 885 889 A (SANOFI SA) 23 December 1998 (1998-12-23)</p> <p>page 3, lines 1-7; examples 1-54</p>	<p>1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56</p>
X	<p>US 5 532 237 A (GALLANT MICHEL ET AL) 2 July 1996 (1996-07-02)</p> <p>column 14, lines 46-66; claim 5; table 1</p>	<p>1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56</p>
P,X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294088 retrieved from XFIRE Database accession no. 9568402, 9579279, 9582786, 9563319</p> <p>abstract &amp; BIOORG. MED. CHEM., vol. 11, no. 18, 2003, pages 3965-3974,</p> <p style="text-align: center;">-/--</p>	<p>1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56</p>

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/US2004/008333

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294089 retrieved from XFIRE Database accession no. 8176641, 8176893  abstract & MOL. PHARMACOL., vol. 48, no. 2, 1995, pages 352-361,	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294090 retrieved from XFIRE Database accession no. 8284677, 8286894, 8287610, 8289960, 8290344,  7780291, 7780333, 8288409, 7231571 abstract & J. MED. CHEM., vol. 42, no. 4, 1999, pages 769-776,	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294091 retrieved from XFIRE Database accession no. 8360417  abstract & J. PHARMACOL., vol. 126, 1999, pages 665-672,	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294092 retrieved from XFIRE Database accession no. 8795090  abstract & J. PHARMACOL. EXP. THER., vol. 284, no. 1, 1998, pages 291-297,	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56

-/--

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/008333

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294093 retrieved from XFIRE Database accession no. 496619</p> <p>abstract &amp; BIOORG. MED. CHEM. LETT., vol. 6, no. 19, 1996, pages 2263-2268,</p>	<p>1-14, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294094 retrieved from XFIRE Database accession no. 2216071</p> <p>abstract &amp; PHARMACOL. TOXICOL., vol. 80, 1997, pages 103-107,</p>	<p>1-14, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294095 retrieved from XFIRE Database accession no. 7226937</p> <p>abstract &amp; BIOORG. MED. CHEM. LETT., vol. 6, no. 1, 1996, pages 17-22,</p>	<p>1-14, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294096 retrieved from XFIRE Database accession no. 497341</p> <p>abstract &amp; XENOBIOTICA, vol. 24, no. 7, 1994, pages 603-612,</p> <p style="text-align: center;">----- -/--</p>	<p>1-14, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56</p>

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/008333

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294097 retrieved from XFIRE Database accession no. 7392775, 7389552, 7390283, 7392387,</p> <p>abstract &amp; J. MDE. CHEM., vol. 41, no. 23, 1998, pages 4521-4532,</p>	<p>1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294098 retrieved from XFIRE Database accession no. 4302866</p> <p>abstract &amp; J. PHARM. PHARMACOL., vol. 53, no. 11, 2001, pages 1525-1532,</p>	<p>1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294099 retrieved from XFIRE Database accession no. 1610624 abstract &amp; CHEM. PHARM. BULL., vol. 42, no. 7, 1994, pages 1459-1462,</p>	<p>1-15,22, 30</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294100 retrieved from XFIRE Database accession no. 4604978 abstract &amp; J. HETEROCYCL. CHEM., vol. 19, 1982, pages 953-956,</p>	<p>1-15,22, 30</p>
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294101 retrieved from XFIRE Database accession no. 5466453 abstract &amp; PHARMAZIE, vol. 47, no. 6, 1992, pages 409-411,</p> <p style="text-align: center;">-/--</p>	<p>1-15,22, 30</p>

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/US2004/008333

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294102 retrieved from Xfire Database accession no. 1904468 abstract & AUST. J. CHEM., vol. 32, 1979, pages 1531-1550, -----	1-15
E	WO 2004/029026 A (EATHERTON ANDREW JOHN ; GIBLIN GERARD MARTIN PAUL (GB); NAYLOR ALAN (G) 8 April 2004 (2004-04-08)  page 13, line 9 - page 15, line 35; claim 1; examples 1-306 -----	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
P,X	FR 2 839 718 A (SANOFI SYNTHELABO) 21 November 2003 (2003-11-21)  Preparation 4.4 on p.22, page 1, lines 1-3; claim 1; example 7 -----	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
P,X	WO 2004/018433 A (EATHERTON ANDREW JOHN ; GIBLIN GERARD MARTIN PAUL (GB); GREEN RICHARD) 4 March 2004 (2004-03-04)  page 17, line 1 - page 19, line 31; claim 1; examples 1-256 ----- -/--	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US2004/008333

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 2004/017920 A (UNIV CONNECTICUT) 4 March 2004 (2004-03-04)	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
	claim 1; tables 1,2	
E	WO 2004/029027 A (EATHERTON ANDREW JOHN ; GIBLIN GERARD MARTIN PAUL (GB); NAYLOR ALAN (G) 8 April 2004 (2004-04-08)	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
	page 2, line 6 - page 3, line 5; claim 1; examples 1,43,45,47,48,50,51,54,56-58,60,63,64,67-6 9,78	
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; XP002294142 retrieved from Xfire Database accession no. 232835 abstract & J. ORG. CHEM., vol. 46, no. 20, 1981, pages 4051-4057,	1-15
X	WO 01/29007 A (LIU QIAN ; MAKRIYANNIS ALEXANDROS (US); UNIV CONNECTICUT (US)) 26 April 2001 (2001-04-26)	1-14,22, 24,26, 28,30, 32,34, 36,38, 40,42, 44,46, 48,50, 52,54,56
	page 18, lines 20,21; claim 1; table 1	

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US2004/008333

## Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 24-29, 32-35, 40, 41, 44-47, 52, 53 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; It is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/008333

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6100259	A	08-08-2000	CA 2278307 A1	23-07-1998
			DE 69822449 D1	22-04-2004
			EP 0971588 A1	19-01-2000
			JP 2001508799 T	03-07-2001
			WO 9831227 A1	23-07-1998
WO 02060447	A	08-08-2002	CA 2436133 A1	08-08-2002
			EP 1363632 A1	26-11-2003
			WO 02060447 A1	08-08-2002
US 5925768	A	20-07-1999	FR 2742148 A1	13-06-1997
			AT 191211 T	15-04-2000
			AU 718763 B2	20-04-2000
			AU 1101097 A	03-07-1997
			BR 9611986 A	17-02-1999
			CA 2239489 A1	19-06-1997
			CN 1207731 A ,B	10-02-1999
			CZ 9801775 A3	14-10-1998
			DE 69607484 D1	04-05-2000
			DE 69607484 T2	26-10-2000
			DK 868420 T3	21-08-2000
			EE 9800171 A	15-12-1998
			EP 0868420 A1	07-10-1998
			EP 0885889 A2	23-12-1998
			ES 2148820 T3	16-10-2000
			WO 9721682 A1	19-06-1997
			GR 3033626 T3	31-10-2000
			HU 9901244 A2	30-08-1999
			JP 2000502080 T	22-02-2000
			JP 3510270 B2	22-03-2004
			NO 982589 A	03-08-1998
			NZ 323962 A	28-05-1999
			PL 327177 A1	23-11-1998
			PT 868420 T	31-08-2000
			RU 2170230 C2	10-07-2001
			SI 868420 T1	31-10-2000
			SK 75898 A3	11-01-1999
			TR 9801041 T2	21-08-1998
			TW 402594 B	21-08-2000
			ZA 9610299 A	18-06-1997
WO 0128557	A	26-04-2001	AU 1095101 A	30-04-2001
			CA 2387138 A1	26-04-2001
			EP 1223929 A1	24-07-2002
			JP 2003512326 T	02-04-2003
			WO 0128557 A1	26-04-2001
WO 0132169	A	10-05-2001	ZA 200203911 A	25-08-2003
			AU 1245801 A	14-05-2001
			CA 2385928 A1	10-05-2001
			EP 1244440 A1	02-10-2002
			HU 0203681 A2	28-04-2003
			JP 2004505006 T	19-02-2004
			NZ 518054 A	31-10-2003
			WO 0132169 A1	10-05-2001
			US 2002173528 A1	21-11-2002
			ZA 200202181 A	18-03-2003

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/008333

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03020217	A	13-03-2003	CA 2457922 A1 EP 1421077 A2 WO 03020217 A2	13-03-2003 26-05-2004 13-03-2003
EP 0885889	A	23-12-1998	FR 2742148 A1 EP 0885889 A2 SI 868420 T1 AT 191211 T AU 718763 B2 AU 1101097 A BR 9611986 A CA 2239489 A1 CN 1207731 A ,B CZ 9801775 A3 DE 69607484 D1 DE 69607484 T2 DK 868420 T3 EE 9800171 A EP 0868420 A1 ES 2148820 T3 WO 9721682 A1 GR 3033626 T3 HU 9901244 A2 JP 2000502080 T JP 3510270 B2 NO 982589 A NZ 323962 A PL 327177 A1 PT 868420 T RU 2170230 C2 SK 75898 A3 TR 9801041 T2 TW 402594 B US 5925768 A ZA 9610299 A	13-06-1997 23-12-1998 31-10-2000 15-04-2000 20-04-2000 03-07-1997 17-02-1999 19-06-1997 10-02-1999 14-10-1998 04-05-2000 26-10-2000 21-08-2000 15-12-1998 07-10-1998 16-10-2000 19-06-1997 31-10-2000 30-08-1999 22-02-2000 22-03-2004 03-08-1998 28-05-1999 23-11-1998 31-08-2000 10-07-2001 11-01-1999 21-08-1998 21-08-2000 20-07-1999 18-06-1997
US 5532237	A	02-07-1996	AU 703913 B2 AU 4616696 A CA 2211836 A1 WO 9625397 A1 EP 0809630 A1 JP 3033076 B2 JP 10508870 T	01-04-1999 04-09-1996 22-08-1996 22-08-1996 03-12-1997 17-04-2000 02-09-1998
WO 2004029026	A	08-04-2004	WO 2004029026 A1	08-04-2004
FR 2839718	A	21-11-2003	FR 2839718 A1 WO 03097597 A2	21-11-2003 27-11-2003
WO 2004018433	A	04-03-2004	WO 2004018433 A1	04-03-2004
WO 2004017920	A	04-03-2004	WO 2004017920 A2 WO 2004017922 A2 US 2004087590 A1	04-03-2004 04-03-2004 06-05-2004
WO 2004029027	A	08-04-2004	WO 2004029027 A1	08-04-2004
WO 0129007	A	26-04-2001	AU 1968901 A	30-04-2001

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/008333

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0129007	A	CA 2387892 A1	26-04-2001
		EP 1224173 A1	24-07-2002
		JP 2003512357 T	02-04-2003
		WO 0129007 A1	26-04-2001

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**